

# Effect and Mechanisms of Medicinal Plants on Dry Eye Disease: A Systematic Review

EZATOLLAH MEMARZADEH<sup>1</sup>, TAHRA LUTHER<sup>2</sup>, SAEID HEIDARI-SOURESHJANI<sup>3</sup>

## ABSTRACT

**Introduction:** Dry Eye Disease (DED) is one of the most common problems and reasons for referring to ophthalmology clinics, which has been exacerbated by the increased use of computers and mobile phones.

**Aim:** The purpose of the present systemic review was to review the effects and mechanism of medicinal plants in DED treatment.

**Materials and Methods:** The key words “Dry eye” or “Keratoconjunctivitis sicca” or “keratitis sicca” in combination with “medicinal plant”, “herb”, and “phyto” were used to conduct the review. Clinical and experimental published articles in English language between 2000-2018 were retrieved from databases including the Institute for Scientific Information (ISI), PubMed and Scopus.

**Results:** Totally, 199 articles were retrieved from the electronic database and finally 17 articles were included in the final analysis. Various plants such as *Buddleja officinalis*, *Aristolelia*

*chilensis*, *Prunus armeniaca*, *Hippophae rhamnoides*, *Lycium barbarum*, and *Rhynchosia volubilis* Lour are effective in treating DED through different mechanisms.

Herbal derivatives such as curcumin, antioxidant glasses, phytoestrogen, ferulic acid, and kaempferol can be used as food supplement independently and in some cases along with chemical drugs. Chinese herbal formulations and compounds such as ‘Chi-Ju-Di-Huang-Wan’, ‘TriphalaGhritaNetratarpan’, and ‘ShengJinRunZaoYangXue’ granules can play a role in inducing anti-oxidant and anti-inflammatory properties in the treatment of the disease.

**Conclusion:** Generally, medicinal plants reduce tear film stability by decreasing osmolarity and increasing tear production. Several mechanisms, including the reduction of Reactive Oxygen Species (ROS) (antioxidant activity), the prevention of cell apoptosis, the modulation of inflammatory factors, and the regulation of androgens, can affect lacrimal glands and membrane cells, thereby helping to treat DED.

**Keywords:** Keratitis sicca, Keratoconjunctivitis sicca, Medicinal herb

## INTRODUCTION

Dry eye disease or keratoconjunctivitis sicca is a multifactorial disease that is developed due to inflammation of the ocular surface and lacrimal glands, meibomian gland dysfunction, and neuropathic pain disorder [1]. Various environmental factors such as air dryness, smoke, allergens, systemic diseases, and age (in premenopausal women) contribute to DED [2-5]. The disease has increased in recent years with the advent of monitors and prolonged use of computers and is one of the most important causes of referral to ophthalmology clinics characterised by symptoms such as scratches and foreign bodies in the eye [6-8].

Due to the high prevalence of DED, it is necessary for physicians to have important information about care, often debilitating symptoms, and preventable and curable nature of the disease [5].

DED treatment is based on a tear substitute. New treatments focus on certain drugs such as secretagogues, lubricants, topical androgens, antibiotics, immunosuppressive drugs, and new anti-inflammatory drugs [9,10]. Meanwhile, herbal treatments have been used in the treatment of various diseases due to being less costly and causing fewer side effects [11-19]. Although, the use of herbal medicines has been used in many countries for the treatment of eye diseases since long time [20-23], however, little information is available on the validity of the efficacy or probable side effects of these drugs. Also, regarding the high prevalence of the disease, and considering that there is still no study in this regard, the authors aimed to investigate the effect and mechanism of medicinal plants in treating DED in the present systemic review.

## MATERIALS AND METHODS


The key words ‘Dry eye’ or ‘Keratoconjunctivitis sicca’ or ‘keratitis sicca’ in combination with medicinal plant, herb, and phyto were used to conduct the review. Relevant articles (from 2000 till date) were retrieved by using keywords from databases including ISI, PubMed, and Scopus with EndNote software. Then, the plants and the plant-based products that were effective on DED were selected and entered to the study according to inclusion criteria. A standard form, which included items such as year of publication, aim and the title of the study, intervention, outcome, variables, journal name, period, and number, was designed. The articles contents that were in accordance with the purpose of the study were recorded in the form and entered to the study with agreement of researchers. The articles whose full texts were not accessible, the articles published in non-English languages, and those that did not address the subject of this review were excluded after the author’s agreement was achieved. [Table/Fig-1] is the flowchart that illustrates how the articles were selected for final analysis.

## RESULTS

The 17 studies that were included in the review and were effective on treatment on DED are tabulated in [Table/Fig-2] [24-40].

## DISCUSSION

Dry eye disease is a multifactorial disease. Hence, the treatment must be based on underlying cause of the disease. Medicinal plants and their derivatives can impact DED in several ways. Medicinal plants improve tear film stability by decreasing osmolarity and inflammation increasing tear production. Tear hyperosmolarity has been reported to begin dry eye inflammation through the



[Table/Fig-1]: Flowchart of the process of selecting and analysing the articles.

activation of epithelial and stromal cells on the ocular surface, which increase the presence of pro-inflammatory cytokines [41]. Inflammation can cause dysfunction in the lacrimal glands cells responsible for tear secretion or retention and leads to ocular surface disease. Plant compounds can be used to regulate levels of gonadal hormones by exhibiting androgen-like activities. As a result, it can inhibit topical inflammation in the lacrimal glands and reduce cell apoptosis [41,42]. In addition, plants and their compounds can contribute to the treatment process by inhibiting the expression of pro-inflammatory cytokines (IL-1 beta, IL-6, and TNF $\alpha$ ) and reducing nitric oxide production [30,43,44]. Artificial tears are the most common treatment for eye diseases therapy; they provide temporary symptomatic relief; however, are unable inhabitation of inflammatory cytokines activity [34]. Also, reduction of ROS (antioxidant activity) and the prevention of cell apoptosis can reduce the symptom of DED. A 8-Hydroxy-2 deoxyguanosine (8-OHdG), 4-Hydroxynonenal (HNE) and Malondialdehyde (MDD) are major biomarkers that cause oxidative damage in ocular surface [41,45]. In DED patients, the levels of these in the tear and conjunctiva increase and cause disruption in the normal functioning of the eye and finally leads to oxidative damage, possibly inflammation and apoptosis [41]. Anti-apoptotic activity and increased expression of some genes in the lacrimal gland epithelial cells represent one of the mechanisms in treating DED [46]. Medicinal plants indicate androgen-like activity and can affect lacrimal glands and membrane cells, thereby contributing to DED treatment. Androgens may increase meibomian gland function and lacrimal gland function [47].

Medicinal plants				
References	Plants	Study Design	Type of administration	Main effects or mechanisms
Peng QH et al., [24]	<i>Buddleja officinalis</i>	Experimental (in vivo)	Extract	Displaying androgen-like activity to keep basic tears secretory volume and tear film stability
Hitoe Set al., [25]	<i>Aristotelia chilensis</i>	Clinical pilot trial	standardised extract (MaquiBright)	Increasing tear fluid volume
Kim CS et al., [26]	<i>Prunus armeniaca</i>	Experimental (in vivo)	Kernel extract	Promoting the secretion of tear fluid and mucin
Kimura Y et al., [27]	<i>Hippophae rhamnoides</i>	Experimental (in vivo)	Pulp oil	Preserving tear secretion and inflammatory cytokines in the lacrimal gland
Chien KJ et al., [28]	<i>Lycium barbarum</i>	Experimental (in vivo)	Extract	Enhancing the tear volume and repairing the damaged ocular surface cells by exhibiting antioxidant and anti-inflammatory activity.
Kang SW et al., [29]	<i>Rhynchosia volubilis Lour.</i>	Experimental (in vivo)	Ethanol extract	Inhibiting squamous metaplasia and apoptosis in the cornea and inhibiting the expression of cytochrome c and Bax, while improving that of Bcl-2.
Phytochemicals				
References	Phytocompound and plant derivatives	Origin	Study Design	Main effects or mechanisms
Chen M et al., [30]	Curcumin	Turmeric ( <i>Curcuma longa</i> )	Experimental (in vivo)	Protecting against hyperosmoticity-induced IL-1 $\beta$ elevation in human corneal epithelial cell via the MAPK pathways and producing anti-inflammatory effect
Scuderi G et al., [31]	Lacrisek (phytoestrogen compound)	$\alpha$ -lipoic acid (100 mg), eicosapentaenoic acid (240 mg), and extract of fenugreek (200 mg), which contains diosgenin (1.3%), steroidal saponins (50%), and alkaloids	Randomised clinical trial	Decreasing tear osmolarity and significantly increasing tear production
Choi W et al., Choi W et al., and Huang JY et al., [32-34]	Antioxidant glasses	Extract of <i>Schizonepetatenuifolia</i> var. <i>japonica</i> Kitagawa, <i>Angelica dahurica</i> Bentham ET hooker, <i>Rehmanniaglutinosa</i> Liboschitz var. <i>purpurea</i> Makino, and <i>Cassia tora</i> L extracts/ <i>Cassiae semen</i> and <i>Ophiopogonis japonicus</i>	Randomised Clinical trial/Experimental (in vivo)	Exerting preventive effect on membrane damage, and inflammation in DED by improving all clinical parameters, IL-1 $\beta$ , IL-6, TNF- $\alpha$ , and IFN- $\gamma$ levels, percentage of CD4+CXCR3+ T cells, goblet cell density, number of 4-HNE-positive cells, and extracellular reactive oxygen species (ROS) production
Chen HC et al., [35]	Ferulic acid and kaempferol	-	Experimental (in vitro)	Ferulic acid and kaempferol lower IL-1 $\beta$ , IL-6, IL-8, and TNF $\alpha$ expression and produce antioxidant effect

Plant-based compounds				
References	Plant-based compounds	Compounds	Study Design	Main effects or mechanisms
Peng QH et al., [36]	Chi-Ju-Di-Huang-Wan	Chi-Ju-Di-Huang-Wan' is the formula 'Liu Wei Di Huang Wan' adds Fructus Lycii and Hos Chrysanthemi	Experimental (in vivo)	Reduce abnormalities in the corneal epithelium and be used as an artificial tear
Chen YL et al., [37].	Chinese herbal treatment	Ji-Sheng-Shen-Qi-Wan, Well Zhen Zi, Dan shen, Gou Qi Zi, Well Zhen Zi, Gu Jing, Hang Bai Ju, and HuaNiu Xi	Case-report	Effects on estrogen receptors and hormonal regulation
Sawant DP et al., [38].	Triphala Ghrita Netratarpan	Phyllanthus emblica, Terminalia chebula, Terminalia bellirica in ghee base	Clinical trial	Reduced the symptoms of the computer vision syndrome
Hu W et al., [39].	Sheng Jin Run ZaoYang Xue granules	Beishashen, Dihuang, Maidong, Shihu, Heshouwu, Gouqi, Juhua, Guizhencao, Zivan and Zhigancao	Clinical trial	Treat DED, which is one of the symptoms of Sjögren's syndrome
Wu SQ et al., [40]	Chinese medicinal plants	Qiwei Baizhu powder, Zhibai Dihuang decoction, Liuwei Dihuang decoction, and kidney-Qi pills	Clinical trial	Reduce levels of interleukin (IL)-1, IL-8, and TNF $\alpha$ and to improve eye dryness

**[Table/Fig-2]:** Medicinal plants, phytochemicals and plant-based compounds effective in dry eye disease [25-41].

Clearly, it should be noted that some herbs did not have any positive effect on DED treatment (in humans or animals) [48,49]. Anticholinergic alkaloids are one of the strongest causes of the disease [50]. The most important plant compounds that lead to the development of DED include niacin, echinacea, and kava, and it should be noted that herbal drugs per se can cause eye complications for consumers, which is due to the natural origin of these products, are usually ignored by the consumers and even by the doctors who prescribe the herbal drugs [51,52].

For example, pouring black tea (*Camellia sinensis*) directly into the cornea with the corneal epithelial defect should be prevented, as it leads to anterior stromal discoloration in individuals [53]. On the other hand, determining the effective dose of medicinal plants is another factor affecting their efficacy. Determining the effective dose in DED treatment means the determination of dose for the highest efficacy and also in the toxic dose [28,35].

In some studies, this issue has been neglected and different quantities of plant extract or derivatives at different intervals have not been studied, which was one of the major problems with these studies [54,55].

## CONCLUSION

Medicinal plants improve tear film stability by decreasing osmolality and increasing tear production. Several mechanisms, including the reduction of ROS (antioxidant activity), the prevention of cell apoptosis, the modulation of inflammatory factors, and the regulation of androgens, can affect lacrimal glands and membrane cells, thereby contributing to DED treatment.

## ACKNOWLEDGEMENTS

The authors would like to acknowledge Research and Technology Deputy of Shahrekord University of Medical Sciences for supporting the present study.

## REFERENCES

- Javadi MA, Feizi S. Dry Eye Syndrome. *J Ophthalmic Vis Res.* 2011;6(3):192-98.
- Gayton JL. Etiology, prevalence, and treatment of dry eye disease. *Clin Ophthalmol.* 2009;3:405-12.
- Hom MM, Nguyen AL, Bielory L. Allergic conjunctivitis and dry eye syndrome. *Ann Allergy Asthma Immunol.* 2012;108(3):163-66.
- Hessen M, Akpek EK. Dry eye: an inflammatory ocular disease. *J Ophthalmic Vis Res.* 2014;9(2):240-50.
- Peck T, Olsakovsky L, Aggarwal S. Dry Eye syndrome in menopause and perimenopausal age group. *J Midlife Health.* 2017;8(2):51-54.
- Logaraj M, Madhupriya V, Hegde S. Computer vision syndrome and associated factors among medical and engineering students in chennai. *Ann Med Health Sci Res.* 2014;4(2):179-85.
- Golden MI, Fries PL. *Dry Eye Syndrome.* Treasure Island (FL): StatPearls Publishing, 2017. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/29262012>
- Dry Eye USA: U.S. National Library of Medicine-The World's Largest Medical Library; 2018 [Available from: <https://www.ncbi.nlm.nih.gov/pubmedhealth/PMHT0023579/>].
- Messmer EM. The pathophysiology, diagnosis, and treatment of dry eye disease. *Dtsch Arztebl Int.* 2015;112(5):71-82.
- Bhavsar AS, Bhavsar SG, Jain SM. A review on recent advances in dry eye: Pathogenesis and management. *Oman J Ophthalmol.* 2011;4(2):50-56.
- Yavangi M, Rabiee S, Nazari S, Farimani-Sanoee M, Amiri I, Bahmanzadeh M, et al. Comparison of the effect of oestrogen plus *Foeniculum vulgare* seed and oestrogen alone on increase in endometrial thickness in infertile women. *J Clin Diagn Res.* 2018;12(1):QC01-QC04.
- Shabanian S, Khallil S, Lorigooini Z, Malekpour A, Heidari-Soureshjani S. The effect of vaginal cream containing ginger in users of clotrimazole vaginal cream on vaginal candidiasis. *J Adv Pharm Technol Res.* 2017;8(2):80-84.
- Solati K, Asadi-Samani M, Heidari-Soureshjani S. Medicinal plants effective on serotonin level: a systematic review. *Br J Pharm Res.* 2017;19(4):01-12.
- Shirani M, Raeisi R, Heidari-Soureshjani S, Asadi-Samani M, Luther T. A review for discovering hepatoprotective herbal drugs with least side effects on kidney. *J Nephropharmacol.* 2017;6(2):38-48.
- Shabanian G, Heidari-Soureshjani S, Rafeiean-Kopaei M, Saadat M, Shabanian M. Therapeutic effects of *Quercus persica* I fruit skin on healing of second-degree burn wounds in animal model. *J Zanjan Uni Med Sci Health Services.* 2017;25(113):81-92.
- Nikfarjam M, Rakhshan R, Ghaderi H. Comparison of effect of *lavandula officinalis* and venlafaxine in treating depression: a double blind clinical trial. *J Clin Diagn Res.* 2017;11(7):KC01-KC04.
- Heidari-Soureshjani S, Asadi-Samani M, Yang Q, Saeedi-Boroujeni A. Phytotherapy of nephrotoxicity-induced by cancer drugs: an updated review. *J Nephropathol.* 2017;6(3):254-63.
- Rafeiean-Kopaei M, Baradaran A, Rafeiean M. Plants antioxidants: From laboratory to clinic. *J Nephropathol.* 2013;2(2):152-53.
- Nasri H. Herbal drugs and new concepts on its use. *J Prev Epidemiol.* 2016;1(1).
- Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol.* 2013;4:177.
- Ukponmwan CU, Momoh N. Incidence and Complications of Traditional Eye Medications in Nigeria in a Teaching Hospital. *Middle East Afr J Ophthalmol.* 2010;17(4):315-19.
- Shenoy R, Bialasiewicz A, Khandekar R, Al Barwani B, Al Belushi H. Traditional medicine in Oman: its role in ophthalmology. *Middle East Afr J Ophthalmol.* 2009;16(2):92-96.
- Wilkinson JT, Fraunfelder FW. Use of herbal medicines and nutritional supplements in ocular disorders: an evidence-based review. *Drugs.* 2011;71(18):2421-34.
- Peng QH, Yao XL, Peng J, Wu QL, Tan HY, Zhang JR. Effects of extract of *buddleja officinalis* eye drops on lacrimal gland cells apoptosis of castrated rabbits with dry eye. *Int J Ophthalmol.* 2010;10(1):40-43.
- Hitoe S, Tanaka J, Shimoda H. MaquiBright standardized maqui berry extract significantly increases tear fluid production and ameliorates dry eye-related symptoms in a clinical pilot trial. *Panminerva Med.* 2014;56(3 Suppl 1):01-06.
- Kim CS, Jo K, Lee IS, Kim J. Topical application of apricot kernel extract improves dry eye symptoms in a unilateral exorbital lacrimal gland excision mouse. *Nutrients.* 2016;8(11): pii:E750.
- Kimura Y, Mori D, Imada T, Izuta Y, Shibuya M, Sakaguchi H, et al. Restoration of tear secretion in a murine dry eye model by oral administration of palmitoleic acid. *Nutrients.* 2017;9(4) pii: E364.
- Chien KJ, Horng CT, Huang YS, Hsieh YH, Wang CJ, Yang JS, et al. Effects of *Lycium barbarum* (goji berry) on dry eye disease in rats. *Mol Med Rep.* 2018;17(1):809-18.
- Kang SW, Kim KA, Lee CH, Yang SJ, Kang TK, Jung JH, et al. A standardized extract of *Rhynchosia volubilis* Lour. exerts a protective effect on benzalkonium chloride-induced mouse dry eye model. *J Ethnopharmacol.* 2018;215:91-100.
- Chen M, Hu DN, Pan Z, Lu CW, Xue CY, Aass I. Curcumin protects against hyperosmoticity-induced IL-1 $\beta$  elevation in human corneal epithelial cell via MAPK pathways. *Exp Eye Res.* 2010;90(3):437-43.
- Scuderi G, Contestabile MT, Gagliano C, Iacovello D, Scuderi L, Avitabile T. Effects of phytoestrogen supplementation in postmenopausal women with dry eye syndrome: a randomized clinical trial. *Can J Ophthalmol.* 2012;47(6):489-92.

- [32] Choi W, Kim JC, Kim WS, Oh HJ, Yang JM, Lee JB, et al. Clinical effect of antioxidant glasses containing extracts of medicinal plants in patients with dry eye disease: a multi-center, prospective, randomized, double-blind, placebo-controlled trial. *PLoS One*. 2015;10(10):e0139761.
- [33] Choi W, Lee JB, Cui L, Li Y, Li Z, Choi JS, et al. Therapeutic efficacy of topically applied antioxidant medicinal plant extracts in a mouse model of experimental dry eye. *Oxid Med Cell Longev*. 2016;2016:4727415.
- [34] Huang JY, Yeh PT, Hou YC. A randomized, double-blind, placebo-controlled study of oral antioxidant supplement therapy in patients with dry eye syndrome. *Clin Ophthalmol*. 2016;10:813-20.
- [35] Chen HC, Chen ZY, Wang TJ, Drew VJ, Tseng CL, Fang HW, et al. Herbal supplement in a buffer for dry eye syndrome treatment. *Int J Mol Sci*. 2017;18(8):pii:E1697.
- [36] Peng Q-H, Yao X-L, Wu Q-L, Tan H-Y, Zhang J-R. Effects of extract of *Buddleja officinalis* eye drops on androgen receptors of lacrimal gland cells of castrated rats with dry eye. *Int J ophthalmol*. 2010;3(1):43.
- [37] Chen Y-L, Hou MC-C. Acupuncture and Chinese herbal treatment for dry eye disease: two case reports. *Med Acupunct*. 2012;24(1):50-55.
- [38] Sawant DP, Parlikar GR, Binorkar SV. Efficacy of Triphala Ghrita Netratarpan in computer vision syndrome. *Int J Res Ayurveda Pharm*. 2013;4(2):244-48.
- [39] Hu W, Qian X, Guo F, Zhang M, Lyu C, Tao J, et al. Traditional Chinese medicine compound ShengJinRunZaoYangXue granules for treatment of primary Sjogren's syndrome: a randomized, double-blind, placebo-controlled clinical trial. *Chin Med J (Engl)*. 2014;127(15):2721-26.
- [40] Wu SQ, Dai Q, Xu JL, Sheng WY, Xu QB, Zhong LY. Combined effect of traditional Chinese and Western medicine on inflammatory factors in patients with diabetes-induced xerophthalmia. *Genet Mol Res*. 2016;15(4): gmr15049030
- [41] Peng QH, Yao XL, Wu QL, Chen M. Effects of extract of *buddleja officinalis* on prevention of dry eye in castrated rabbits. *Zhonghua Yan Ke Za Zhi*. 2008;44(11):1011-19.
- [42] Yao XL, Peng QH, Chen QL, Tang YH, Zhong Q. Effects of total flavonoid of chrysanthemum on Fas and FasL expression in lacrimal gland cells of male castrated rabbits with dry eye. *Int Eye Sci*. 2014;14(10):1749-54.
- [43] Wozniak A, Paduch R. Aloe vera extract activity on human corneal cells. *Pharm Biol*. 2012;50(2):147-54.
- [44] Paduch R, Wozniak A, Niedziela P, Rejdak R. Assessment of Eyebright (*Euphrasia Officinalis* L.) Extract activity in relation to human corneal cells using in vitro tests. *Balkan Med J*. 2014;31(1):29-36.
- [45] Barrera G. Oxidative stress and lipid peroxidation products in cancer progression and therapy. *ISRN Oncol*. 2012;2012:137289.
- [46] Wang F, Peng QH, Li HZ, Wang F, Yao XL, Li WJ. Influence for Cell model of dry eye after intervention of plasma containing *Buddleja Officinalis* Flavone and the expression of Bax mRNA and Bcl-2 mRNA in Lacrimal gland epithelial cells. *Int Eye Sci*. 2012;12(10):1836-40.
- [47] Azcarate PM, Venincasa VD, Feuer W, Stanczyk F, Schally AV, Galor A. Androgen Deficiency and Dry Eye Syndrome in the Aging Male. *Invest Ophth Vis Sci*. 2014;55(8):5046-53.
- [48] Abokyi S, Koffuor GA, Ofori EA, Thomford AK, Osei KA. Evaluating the effect of an aqueous extract of *Pistia stratiotes* Linn (Araceae) on tear secretion and tear film stability in ICR mice. *J Appl Pharm Sci*. 2014;4(8):1-5.
- [49] Chan CC, Lau WN, Chiu SP, Chen LC, Choi WK, Tang GW. A pilot study on the effects of a Chinese herbal preparation on menopausal symptoms. *Gynecol Endocrinol*. 2006;22(2):70-73.
- [50] Askeroglu U, Alleyne B, Guyuron B. Pharmaceutical and herbal products that may contribute to dry eyes. *Plast Reconstr Surg*. 2013;131(1):159-67.
- [51] Fraunfelder FW. Ocular side effects from herbal medicines and nutritional supplements. *Am J Ophthalmol*. 2004;138(4):639-47.
- [52] Fraunfelder FW. Ocular side effects associated with dietary supplements and herbal medicines. *Drugs Today*. 2005;41(8):537-45.
- [53] Achiron A, Birger Y, Karmona L, Avizemer H, Bartov E, Rahamim Y, et al. Corneal staining and hot black tea compresses. *Isr Med Assoc J*. 2017;19(3):152-55.
- [54] Jafari T. Antioxidants; helpful or harmful? *Annals of Research in Antioxidants*. 2016;1(2):e13.
- [55] Baradaran A. Administration of herbal drugs in geriatric individuals; trends on its helps and hazards. *Geriatrics Persia*. 2017;1(1):e01.

**PARTICULARS OF CONTRIBUTORS:**

1. Assistant Professor, Department of Ophthalmology, Shahrekord University of Medical Sciences, Shahrekord, Iran.
2. Director, Department of General Surgery, University of Michigan, Ann Arbor, MI, USA.
3. MSc, Deputy of Research and Technology, Shahrekord University of Medical Sciences, Shahrekord, Iran.

**NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:**

Dr. Saeid Heidari-Soureshjani,  
Deputy of Research and Technology, Shahrekord University of Medical Sciences, Shahrekord, Chahrmahal va Bakhtiari, Iran.  
E-mail: heidari\_62@yahoo.com

Date of Submission: **Mar 13, 2018**  
Date of Peer Review: **May 23, 2018**  
Date of Acceptance: **Jun 15, 2018**  
Date of Publishing: **Sep 01, 2018**

**FINANCIAL OR OTHER COMPETING INTERESTS:** None.