

Evaluation of Usefulness of Mannheim Peritonitis Index and APACHE II Score in Predicting Mortality and Morbidity in Patients with Peritonitis- A Prospective Diagnostic Test Study

D PAUL TRINITY STEPHEN¹, VIJAY ABRAHAM², REKA KARUPPUSAMI³

ABSTRACT

Introduction: Peritonitis, defined as inflammation of the peritoneal cavity can be of various causes, and is one of the most common surgical emergencies. This continues to be a challenge to diagnose and treat. Early intervention is essential to select patients who will need intensive care which brings out better outcome for the patients. This also helps us use the resources optimally. Over years, many scoring systems have been developed and studied to predict outcomes in patients with peritonitis.

Aim: To evaluate the ability of Mannheim Peritonitis Index (MPI) and APACHE II (Acute Physiology And Chronic Health Evaluation II) scores in predicting mortality and morbidity in patients with peritonitis.

Materials and Methods: A prospective, observational study was conducted at Christian Medical College and Hospital, Vellore, Tamil Nadu, India, for a period of two years from September 2014 to August 2016. A total of 78 patients were recruited for this study. These patients were scored with MPI and APACHE II scores. The primary outcome studied was in hospital death or discharge. The secondary outcome studied was morbidity in terms of local and systemic complications. The risk factors associated with mortality in patients with peritonitis were also

studied. The best cut-off value for MPI and APACHE II from the data was calculated using Yuden index. The sensitivity, specificity and likelihood ratios were calculated and presented with 95% CI. The sub-group analysis was done for risk factors and complications.

Results: There were more males than females. Age ≥ 48 years ($p=0.002$) and Serum creatinine ≥ 1.3 g/dL ($p=0.012$) were found to be significant risk factors for mortality. The sensitivity and specificity of MPI ≥ 27 in predicting mortality was found to be 90% and 57% respectively ($p=0.006$). The sensitivity and specificity of APACHE II score ≥ 10 in predicting mortality was found to be 40% and 78%, respectively ($p=0.47$). MPI scores ≥ 27 were strongly associated with morbidity like prolonged ICU stay ($p=0.004$), mechanical ventilation requirement ($p=0.001$) and need for dialysis ($p=0.035$).

Conclusion: Present study showed MPI to be a better predictor of mortality than APACHE II, though APACHE II showed better specificity. MPI score also was helpful in predicting morbidity such as prolonged ICU stay, mechanical ventilation requirement postoperatively and need for dialysis postoperatively. MPI was easier to use as it contained lesser variables. MPI could be of use in rural areas with no facility for laboratory investigations and blood gas analysis.

Keywords: Outcomes in peritonitis, Peritonitis scoring systems, Risk factors

INTRODUCTION

Peritonitis is defined as inflammation of the peritoneum due to localised or generalised infections. Peritonitis is one of the most common infections and an important problem that a surgeon encounters. Despite the surgical treatment, intensive care treatment and advance in antibiotic therapy and a good understanding of the patho-physiology, the mortality rates of perforation peritonitis are still high ranging from 5.6%-56% [1-4]. Hence, early prognostic evaluation of abdominal sepsis is preferred to select high-risk patients for more aggressive therapeutic procedures and to classify the severity of the disease. Treatment of peritonitis is primarily surgical and early intervention is always preferred [1,5]. Different scoring systems have been used to predict the outcome of patients with peritonitis. These scores can be a good tool to predict the prioritisation of treatment and care of patients and also help to predict the prognostic factors that affect morbidity and mortality in patients with peritonitis. Many scores have been developed and studied over years which include MPI, APACHE II score, POSSUM (Physiological and Operative Severity Score for the enumeration of Mortality and Morbidity) [6-8], Sequential Organ Failure Assessment (SOFA) [9,10] and more.

MPI score was developed by Linder MM et al., [3]. It was based on the retrospective analysis of the data from patients with peritonitis. The MPI is a specific score which has a very good accuracy and allows the determination of the individual prognosis of patients with peritonitis. APACHE II score was developed by Knaus WA [11]. It was devised to stratify prognosis in a group of ill patients and for the determination of success of treatment. Billing A et al., assessed the validity of MPI and its predictive power in 2003 patients and concluded it is an easy and reliable means for risk evaluation in patients with peritonitis [12]. Demmel N et al., in a study of 438 patients, showed MPI >26 showed sensitivity and specificity of 88% and 78%, respectively [13]. Kulkarni SV et al., studied APACHE II scoring system in perforation peritonitis and concluded it was 100% specific at higher scores in predicting mortality [14]. Studies done to compare MPI and APACHE II in predicting outcome showed both scores had comparable sensitivity and specificity but APACHE II was more accurate [15,16]. Malik A et al., in a study done to compare MPI and APACHE II scores in predicting mortality in patients with peritonitis, said both scores showed significance in predicting outcome [17].

As the available literature was not clearly in favour of one score (though APACHE was favoured by many), we wanted to assess

the ability of MPI and APACHE II score to predict outcome in the population. MPI is simple and easy. If MPI could be proven to be as good as APACHE II score, then this could be of good use in rural hospitals (secondary hospitals associated with our centre) lacking facilities for blood gas and other investigations required to use complex scoring systems like APACHE II score.

The aim was to study the ability of MPI and APACHE II score in predicting outcome in patients with peritonitis.

MATERIALS AND METHODS

This prospective observational study was conducted in the Department of General Surgery at Christian Medical College and Hospital, Vellore, Tamil Nadu, India, over a period of two years from September 2014 to August 2016. Ethical clearance was obtained from the institutional research board with study number 9049. Sample size calculation was done based on the sensitivity of MPI score, as in earlier studies, which was found to be about 90% [12,13,18]. With a precision of 15% and 95% desired confidence interval, the number of deaths to be studied was found to be 15. In order to study 15 deaths, with incidence of mortality being 20% in patients with peritonitis, we need to study 75 patients. We finally included 78 patients in the study. All patients more than 16 years of age with secondary peritonitis were included in the study. Patients with peritonitis secondary to trauma, pancreatitis, spontaneous bacterial peritonitis or primary peritonitis were excluded from the study.

A total of 78 patients were recruited. All patients had routine preoperative blood work up done and arterial blood gas was also done. Chest X-ray erect and abdominal X-rays erect and supine were done in all cases. All patients were resuscitated with intravenous fluids and were empirically started on broad spectrum antibiotics at presentation. Nasogastric tube decompression was also done and patients were diagnosed to be peritonitis if they had examination findings of tenderness with guarding or rigidity. All patients underwent emergency laparotomy and surgical procedure done was of the surgeon's choice. All investigations and surgical procedures were carried out with proper informed and written consent as appropriately.

The patients were scored with two scoring systems at admission or within 24 hours. If blood parameters were tested more than once, the most deranged value was used for scoring. The two scores used were MPI and APACHE II score. MPI score used eight risk factors which were found to be significantly associated with prognosis in patients with peritonitis [3]. The maximum score was 47. The best cut-off points used in various studies included 21, 26 and 29 [19-21]. In present study, it was decided to find a new cut-off value for each score and use that to assess the outcome. This was done using Yuden index as described in the statistical analysis. APACHE II score had 2 parts. The first one dealt with acute physiology and the second with chronic health evaluation. It was primarily designed for intensive care unit. It utilises 12 values and determines the outcome [11]. Patients were followed-up to discharge and the primary outcome assessed was in-hospital death or discharge. Secondary outcomes assessed included morbidity and risk factors for mortality in peritonitis. Morbidity was studied in terms of local and systemic complications. The local complications studied were wound infection, wound dehiscence, intra-abdominal collection, anastomotic leak and reoperation. The systemic complications studied were patients requiring dialysis for more than 48 hours postoperative, patients requiring mechanical ventilation for more than 48 hours postoperatively, patients with Glasgow Coma Scale (GCS) less than 8 despite withholding sedation for more than 48 hours, mean hospital stay and mean ICU stay.

STATISTICAL ANALYSIS

Data was entered in EPIDATA software. For continuous data, the descriptive statistics Mean±SD and for non-normally distributed

interval data and ordinal data, median (IQR) was presented. Number of patients and percentage was presented for categorical data. The Chi-square test, Fisher's-exact test (when expected count is less for cell) and Yates continuity correction (where the cell have zero count) was used to find association between two categorical variables. Mann-Whitney U test was performed to compare two groups-ICU stay and hospital stay. The best cut-off value for MPI and APACHE II for present study data was calculated using Yuden index. Receiver Operator Characteristic (ROC) analysis was performed for MPI and APACHE II score. The sensitivity, specificity and likelihood ratios were calculated. All tests were two-sided at $\alpha=0.05$ level of significance. The Statistical Analysis Software (SAS) package (SAS® Institute Inc., USA, version 9.2) was used for statistical evaluation.

RESULTS

Of the 78 patients studied, 59 (75.6%) were males and 19 (24.4%) were females ($p=0.65$) [Table/Fig-1]. The mean age was 48.45 ± 16.67 years. The lowest age was 18 years and the oldest patient was 85-year-old. The minimum heart rate was found to be 68 beats/minute and maximum of 160/minute. The minimum and maximum duration of symptoms were 1 day and 14 days, respectively. The minimum

Patient characteristics	Outcome		Total N=78 N (%)	p-value
	Non-survivors (n=10) N (%)	Survivors (n=68) N (%)		
Sex†				
Male	7 (9%)	52 (66.7%)	59 (75.7%)	0.651
Female	3 (3.8%)	16 (20.5%)	19 (24.3%)	
Age (years)‡				
≤48	0 (0%)	36 (46.15%)	36 (46.15%)	0.002
>48	10 (12.82%)	32 (41.02%)	42 (53.85%)	
Diabetes‡				
Yes	1 (1.3%)	6 (7.7%)	7 (9%)	0.902
No	9 (11.5%)	62 (79.5%)	71 (91%)	
Hypertension‡				
Yes	0 (0%)	8 (10.26%)	8 (10.26%)	0.252
No	10 (12.82%)	60 (76.92%)	70 (89.74%)	
Duration of symptoms‡				
<2 days	4 (5.1%)	23 (29.4%)	27 (34.6%)	0.701
≥2 days	6 (7.7%)	45 (57.7%)	51 (65.4%)	
ASA score‡				
ASA 1	6 (7.6%)	49 (62.8%)	55 (70.5%)	0.430
ASA 2	4 (5.1%)	19 (24.4%)	23 (29.5%)	
Heart rate (beats/minute)				
≤110	5 (6.4%)	34 (43.5%)	39 (50%)	1.001
>110	5 (6.4%)	34 (43.5%)	39 (50%)	
Systolic BP (mmHg)‡				
≥100	2 (2.5%)	25 (32.05%)	27 (34.6%)	0.292
<100	8 (10.3%)	43 (55.12%)	51 (65.4%)	
Total counts (per cubic mm)‡				
4000-12000	3 (3.8%)	32 (41.02%)	35 (44.9%)	0.301
<4000&>12000	7 (9%)	36 (46.15%)	43 (55.1%)	
Left shift‡				
Yes	8 (10.3%)	58 (74.4%)	66 (84.7%)	0.663
No	2 (2.5%)	10 (12.8%)	12 (15.3%)	
Creatinine (g/dL)‡				
≤1.3	4 (5.1%)	53 (68%)	57 (73.1%)	0.012
>1.3	6 (7.6%)	15 (19.2%)	21 (26.9%)	

[Table/Fig-1]: Patient characteristics vs Outcome.

p-value was obtained from the Chi-square test, †Fisher's-exact test (when expected count was less for cell); ‡Yates continuity correction (where the cells have zero count); ASA-American society of anesthesiologist


and maximum systolic blood pressure was found to be 70 mm Hg and 200 mm Hg, respectively. The lowest leukocyte counts noted was 1100/cubic mm and highest counts were 82000/cubic mm. The lowest and highest creatinine values noted were 0.36 g/dL and 6.35 g/dL, respectively.

Majority of the patients (47.4%) were found to have peritonitis secondary to duodenal perforation. The second commonest cause was due to small bowel perforation. Gall bladder perforation was the least common cause for peritonitis among the patients [Table/Fig-2].

Site of perforation	Number (percentage)
Duodenal and pre-pyloric perforation	37 (47.4%)
Ileal (small bowel perforation)	16 (20.5%)
Appendix	14 (17.9%)
Colon	6 (7.6%)
Stomach	3 (3.8%)
Gall bladder	2 (2.5%)

[Table/Fig-2]: Aetiology of perforation.

The best cut-off values for MPI and APACHE II scores were calculated using Yuden index and were found to be 27 and 10 respectively. The area under the curve was found to be 0.78 and 0.66 for MPI and APACHE II scores, respectively [Table/Fig-3].



[Table/Fig-3]: Receiver Operator Characteristic analysis (ROC) for MPI and APACHE II score.

The over-all mortality in present study was 10 patients which was 13%. Among the 10 mortalities, nine patients (90%) had an MPI score of ≥ 27 (p -value=0.005). Out of the 10 non-survivors, four had an APACHE II score of ≥ 10 (p -value=0.435). The mean MPI and APACHE II scores among the non-survivors were found to be 32.5 ± 4.8 and 10.6 ± 3.6 , respectively.

The sensitivity, specificity, positive predictive value, negative predictive value and accuracy rate with a cut-off of 10 for APACHE II, was found to be 40%, 78%, 21%, 89% and 67.5%, respectively. The sensitivity, specificity, positive predictive value, negative predictive values and accuracy rate for MPI score with a cut-off of 27, was found to be 90%, 57%, 23%, 97.5% and 61.5%, respectively.

MPI score ≥ 27 was able to predict need for dialysis ($p=0.035$), need for mechanical ventilation ($p=0.001$) and median ICU stay ($p=0.004$). APACHE II ≥ 10 was unable to predict any morbidity factors that we studied [Table/Fig-4,5].

DISCUSSION

Peritonitis is a common surgical emergency encountered. The mortality and morbidity in such patients have been studied to be high even up to 60%. The mortality rate in present study was found to be 13% which was comparable with most studies [2,22,23].

	MPI ≥ 27 N (%)	MPI < 27 N (%)	p-value
Wound infection [†]	7 (8.97%)	9 (11.53%)	0.651
Wound dehiscence [†]	3 (3.8%)	2 (2.5%)	0.602
Re-operation [†]	2 (2.5%)	1 (1.3%)	0.523
Intra-abdominal collection [†]	5 (6.4%)	6 (7.6%)	0.801
Anastomotic leak	0 (0%)	1 (1.3%)	0.320
Dialysis	4 (5.12%)	0 (0%)	0.035
Mechanical ventilation > 48 hours [†]	25 (32.05%)	11 (14.1%)	0.001
GCS < 8	1 (1.3%)	0 (0%)	0.304
ICU stay(days) ^{††} , Median (IQR)	3 (0,7)	0 (0,3)	0.004
Hospital stay(days) ^{††} , Median(IQR)	8 (7,12)	10 (7,13)	0.306

[Table/Fig-4]: MPI vs. Morbidity.

p-value is obtained from the [†]Fischer's-exact test (when expected count is less for cell) and Yates continuity correction (where the cells have zero count); ^{††}Values are presented as Median (IQR); Median (25th percentile, 75th percentile) and p-value is obtained from nonparametric Mann-Whitney U test

	APACHE ≥ 10 N (%)	APACHE < 10 N (%)	p-value
Wound infection [†]	5 (6.4%)	11 (14.1%)	0.862
Wound dehiscence [†]	1 (1.2%)	4 (5.12%)	0.602
Re-operation [†]	0 (0%)	3 (3.8%)	0.254
Intra-abdominal collection [†]	4 (5.12%)	7 (8.97%)	0.561
Anastomotic leak	0 (0%)	1 (1.3%)	0.515
Dialysis	2 (2.5%)	2 (2.5%)	0.353
Mechanical ventilation > 48 hours [†]	13 (16.6%)	23 (29.48%)	0.235
GCS < 8 [†]	0 (0%)	1 (1.3%)	0.515
ICU stay (days) ^{††} , Median (IQR)	3 (0,8)	0 (0,4)	0.073
Hospital stay (days) ^{††} , Median (IQR)	8 (6,13)	9 (7,12)	0.673

[Table/Fig-5]: APACHE II vs. Morbidity.

Values are presented as number (percentage); P value is obtained from the [†]Fischer's exact test (when expected count is less for cell) and Yates continuity correction (where the cells have zero count); ^{††}Values are presented as Median (IQR); Median (25th percentile, 75th percentile) and p-value is obtained from nonparametric Mann-Whitney U test

The outcome in such patients depends on several factors like age, duration of symptoms, co-morbidities and more [22,24,25]. This has been studied by many studies before and most of them concluded increasing age, longer duration of symptoms and organ failure to be significant risk factors to predict mortality. In present study, we found only age ≥ 48 years and Serum creatinine ≥ 1.3 g/dL to be significant risk factors. Other factors such as co-morbidities, duration of symptoms, American Society of Anesthesiologists (ASA) score, tachycardia, hypotension, abnormal white cell counts, and left shift were not found to be independent risk factors in predicting mortality in peritonitis as compared to previous studies [22,24,25].

In present study, majority of patients were males (75.6%) compared to females (24.3%) as supported by other studies [15]. The aetiology of perforation in present study was commonly due to duodenal perforation as in previously described studies [26-28]. The mean MPI and APACHE II scores in non-survivors in present study were comparable to prior studies [16,29,30]. The sensitivity, specificity, positive and negative predictive values and accuracy rates of MPI and APACHE II as compared with other studies is described in [Table/Fig-6,7] [3,10,12,16,31-34]. We found comparable accuracy rates of MPI and APACHE II from previous studies [16,35].

MPI was found to be more sensitive than APACHE II in present study, though studies prior studies showed better APACHE II sensitivity [14,17]. Many authors have reported APACHE II to have better prognostic power for outcome prediction than MPI as it includes physiological variables [4,11,14]. In present study, the accuracy rate of APACHE II (67.5%) was higher than that of MPI (61.5%), as described in Dino H et al., [10]. In present study, we inferred that MPI was found to have better sensitivity; however the specificity and accuracy rates were better with APACHE II score. MPI on the other

Study	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Accuracy rate
Billing A et al., [12]	76%	58%	-	-	-
Valliant TAL et al., [31]	87%	88%	93%	94%	-
Linder MM et al., [3]	88%	90%	87%	90%	-
Dani DT et al., [32]	90.6%	91.7%	67.44%	98.12%	-
Ojuka A et al., [33]	84.2%	90.7%	75.9%	94.2%	-
Kumar P et al., [16]	100%	91%	69%	100%	69%
Present study	90%	57%	23%	97.5%	61.5%

[Table/Fig-6]: Sensitivity, specificity, positive and negative predictive value and accuracy rate for MPI score [3,12,16,31-33].

Study	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Accuracy rate
Dino H et al., [10]	82.5%	55.2%	54.7%	82.8%	66%
Headly J et al., [34]	54%	87%	-	-	-
Kumar P et al., [16]	85%	100%	100%	96%	83.3%
Present study	40%	78%	21%	89%	67.5%

[Table/Fig-7]: Sensitivity, specificity, positive and negative predictive value and accuracy rate for APACHE II score [10,16,34].

hand, was able to predict need for dialysis mechanical ventilation and prolonged ICU stay. This partly correlated with a study done previously by Gupta N et al., though the study also showed APACHE II to be good predictor of some complications such as need for ventilator stay and longer hospital stay [36]. As in present study, they also found both the scores to be poor predictors of complications like wound infection and anastomotic leak. Adesunkanmi ARK et al., in 69 patients with peritonitis, did not find any significantly higher APACHE II scores in those patients who survived with postoperative complications and had long duration of hospital stay [37]. Similarly, in present study APACHE II was not a good predictor of morbidity.

Limitation(s)

We consider the small sample size a limitation to present study results. Though we aimed to find at least 15 deaths in studying the sample size, we were able to study only 10 deaths. This could contribute to our result of showing the two scores to be not very accurate (both had <70% accuracy rate). It would be prudent to study a larger population and assess the scoring systems.

CONCLUSION(S)

MPI is a simple tool as compared to APACHE II which requires laboratory investigations including blood gas analysis. MPI also takes into consideration the aetiology of peritonitis and the nature of contamination unlike APACHE II. On the other hand, it fails to include the underlying physiological derangements of the patient which is necessary to prognosticate patients requiring intensive care. MPI proves to be a good prognosticative indicator from present study and we recommend it as an alternative tool for usage especially in areas with poor infra-structure like rural hospitals where blood gas analysis would not be possible. However, MPI also needs operative findings to complete the score, hence cannot be used preoperatively. APACHE II continues to be more specific and accurate in predicting patients with mortality.

REFERENCES

[1] Giesslering U, Petersen S, Freitag M, Kleine-Kraneburg H, Ludwig K. Surgical management of severe peritonitis. *Zentralbl Chir.* 2002;127(7):594-97.

[2] Qureshi AM, Zafar A, Saeed K, Qudus A. Predictive power of Mannheim Peritonitis Index. *J Coll Physicians Surg Pak.* 2005;15(11):693-96.

[3] Linder MM, Wacha H, Feldmann U, Wesch G, Streifensand RA, Gundlach E. The Mannheim peritonitis index. An instrument for the intraoperative prognosis of peritonitis. *Chirurg.* 1987;58(2):84-92.

[4] Schein M, Gecelter G, Freinkel Z, Gerding H. APACHE II in emergency operations for perforated ulcers. *Am J Surg.* 1990;159(3):309-13.

[5] Farthmann EH, Schöffel U. Principles and limitations of operative management of intraabdominal infections. *World J Surg.* 1990;14(2):210-17.

[6] Campillo-Soto A, Flores-Pastor B, Soria-Aledo V, Candel-Arenas M, Andrés-García B, Martín-Lorenzo JG, et al. The POSSUM scoring system: An instrument for measuring quality in surgical patients. *Cir Esp.* 2006;80(6):395-99.

[7] Chatterjee AS, Renganathan DN. POSSUM: A scoring system for perforative peritonitis. *J Clin Diagn Res.* 2015;9(4):PC05-09.

[8] Vishwani A, Gaikwad VV, Kulkarni RM, Murchite S. Efficacy of POSSUM Scoring system in predicting mortality and morbidity in patients of peritonitis undergoing laparotomy. *Int J Sci Stud.* 2014;2(4):29-36 (2020 Jun 19).

[9] Vincent JL, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. *Intensive Care Med.* 1996;22(7):707-10.

[10] Dino H, Murni T, Redjeki I. Comparison of Apache II, SOFA, and Modified SOFA Scores in predicting mortality of surgical patients in intensive care unit at Dr. Hasan Sadikin General Hospital. *Critical Care and Shock.* 2009;12.

[11] Knaus WA. APACHE 1978-2001: The development of a quality assurance system based on prognosis: Milestones and personal reflections. *Arch Surg.* 2002;137(1):37-41.

[12] Billing A, Fröhlich D, Schildberg FW. Prediction of outcome using the Mannheim peritonitis index in 2003 patients. *Peritonitis Study Group. Br J Surg.* 1994;81(2):209-13.

[13] Demmel N, Muth G, Maag K, Osterholzer G. Prognostic scores in peritonitis: The Mannheim Peritonitis Index or APACHE II?. *Langenbecks Arch Chir.* 1994;379(6):347-52.

[14] Kulkarni SV, Naik AS, Subramanian N. APACHE-II scoring system in perforative peritonitis. *Am J Surg.* 2007;194(4):549-52.

[15] Mishra A, Singh KK, Jain V. A comparative analysis between Mannheim peritonitis score and acute physiological and chronic health evaluation II score in predicting prognosis of patients of perforation peritonitis. *International Journal of Research in Medical Sciences.* 2019;8(1):10-14.

[16] Kumar P, Singh K, Kumar A. A comparative study between Mannheim peritonitis index and APACHE II in predicting the outcome in patients of peritonitis due to hollow viscous perforation. *International Surgery Journal.* 2017;4(2):690-96.

[17] Malik A, Wani K, Dar L, Wani M, Wani R. Mannheim Peritonitis Index and APACHE II- Prediction of outcome in patients with peritonitis. *Turkish Journal of Trauma & Emergency Surgery: TJTES.* 2010;16:27-32.

[18] Notash AY, Salimi J, Rahimian H. Evaluation of Mannheim peritonitis index and multiple organ failure score in patients with peritonitis. *Indian J Gastroenterol.* 2005;24(5):197-200.

[19] Neri A, Marrelli D, Scheiterle M, Di Mare G, Sforza S, Roviello F. Re-evaluation of Mannheim prognostic index in perforative peritonitis: Prognostic role of advanced age. A prospective cohort study. *International Journal of Surgery.* 2015;13:54-59.

[20] Kusumoto Y, Nakagawa M, Watanabe A, Ishikawa H, Sakaguchi T, Yamada T, et al. Study of Mannheim Peritonitis index to predict outcome of patients with peritonitis. *The Japanese Journal of Gastroenterological Surgery.* 2004;37:07-13.

[21] Biondo S, Ramos E, Fraccalvieri D, Kreisler E, Ragué JM, Jaurieta E. Comparative study of left colonic peritonitis severity score and manheim peritonitis index. *BJS (British Journal of Surgery).* 2006;93(5):616-22.

[22] Singh R, Kumar N, Bhattacharya A, Vajifdar H. Preoperative predictors of mortality in adult patients with perforation peritonitis. *Indian J Crit Care Med.* 2011;15(3):157-63.

[23] Muralidhar VA, Madhu CP, Sudhir S, Srinivasarangan M. Efficacy of Mannheim Peritonitis Index (MPI) score in patients with secondary peritonitis. *J Clin Diagn Res.* 2014;8(12):NC01-03.

[24] Khan PS, Dar LA, Hayat H. Predictors of mortality and morbidity in peritonitis in a developing country. *Ulus Cerrahi Derg.* 2013;29(3):124-30.

[25] Lamme B, Mahler CW, van Ruler O, Gouma DJ, Reitsma JB, Boermeester MA. Clinical predictors of ongoing infection in secondary peritonitis: Systematic review. *World J Surg.* 2006;30(12):2170-81.

[26] Arasu VT, Lakshminpathy N. A prospective study of evaluation of manheim peritonitis index to predict outcome of patients with peritonitis. *International Journal of Contemporary Medical Research.* 2016;3(11):3339-41.

[27] Koppad SN, Vandakudri AB, Desai M, Kodliwadmath H. Analysis of manheim peritonitis index scoring in predicting outcome in patients with peritonitis secondary to hollow viscous perforation. *International Surgery Journal.* 2016;3(3):1116-20.

[28] Reddy MV, Reddy TA, Satyadev, Teja BR, ShanmugaRaju P. Application of APACHE II Score in assessing the severity and outcome in peritonitis due to hollow viscous perforation. *International Surgery Journal.* 2019;6(3):940-43.

[29] Sharma VK, Basnet RB. Evaluation of predictive power of manheim peritonitis index. *Post-Graduate Medical Journal of NAMS.* 2010;10(02).

[30] Naveen P, Dhannur PK. Modified APACHE II scoring and Mannheims peritonitis Index (MPI) in predicting the outcome of patients with peritonitis secondary to hollow viscous perforation. *Int J Surg Sci.* 2019;3(3):403-07.

[31] Vaillant TAL, Morejón CS, Barbeito TOT. Scoring systems in sepsis. *Rev Cub Med Mil.* 2012;41(4):394-406.

[32] Dani DT, Ramachandra PL, Nair DR, Sharma DD. Evaluation of prognosis in patients' with perforation peritonitis using manheim's peritonitis index. *International Journal of Scientific and Research Publications.* 2015;5(5).

- [33] Ojuka A, Ekwaro L, Kakande I. Causes and patterns of peritonitis at St. Francis Hospital Nsambya, Kampala-Uganda. *East Cent Afr J Surg*.2014;19(3):99-106.
- [34] Headley J, Theriault R, Smith TL. Independent validation of APACHE II severity of illness score for predicting mortality in patients with breast cancer admitted to the intensive care unit. *Cancer*. 1992;70(2):497-503.
- [35] Ohmann C, Wittmann DH, Wacha H. Prospective evaluation of prognostic scoring systems in peritonitis. *Peritonitis Study Group. Eur J Surg*. 1993;159(5):267-74.
- [36] Gupta N, Agrawal H, Gupta AK, Naskar D, Durga CK. A comparative study between APACHE II scoring and Mannheim Peritonitis Index to assess prognosis in perforation peritonitis- SAGES ARCHIVES. 2020 Jun 19.
- [37] Adesunkanmi ARK, Oseni SA, Adejuyigbe O, Agbakwuru EA. Acute generalised peritonitis in African children: Assessment of severity of illness using modified APACHE II score. *ANZ Journal of Surgery*. 2003;73(5):275-79.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of General Surgery, Christian Medical College, Vellore, Tamil Nadu, India.
2. Senior Medical Practitioner, Department of Upper GI Surgery, The Queen Elizabeth Hospital, Woodville South, South Australia, Australia.
3. Lecturer, Department of Biostatistics, Christian Medical College, Vellore, Tamil Nadu, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. D Paul Trinity Stephen,
Surgery 4 Office, Paul Brand Building, Christian Medical Collge and Hospital,
Vellore-632004, Tamil Nadu, India.

E-mail: pl.trinity@gmail.com

PLAGIARISM CHECKING METHODS: [\[Jain H et al.\]](#)

- Plagiarism X-checker: Jun 30, 2020
- Manual Googling: Sep 05, 2020
- iThenticate Software: Sep 22, 2020 (14%)

ETYMOLOGY: Author Origin**AUTHOR DECLARATION:**

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. NA

Date of Submission: **Jun 29, 2020**

Date of Peer Review: **Jul 15, 2020**

Date of Acceptance: **Sep 07, 2020**

Date of Publishing: **Oct 01, 2020**