

# A Cross-sectional Study of Anatomical Variations of Coronary Ostia in the Adult Human Hearts and its Clinical Significance

NAJMA MOBIN<sup>1</sup>, H BASAVANAGOWDAPPA<sup>2</sup>, B MADHU<sup>3</sup>

## ABSTRACT

**Introduction:** The knowledge of anatomical variations of the coronary ostia is an important factor in the study of Coronary Heart Disease (CHD). With the adaptation of sedentary life style and increasing stress, CHD has become the major cause of death in many countries.

**Aim:** To describe the location, size and shape of the coronary ostia in cadaveric hearts.

**Materials and Methods:** Descriptive cross-sectional study was carried out in 110 adult human hearts, 90 males and 20 females with a mean age of 66 years, which were collected from the mortuary from March 2017 to March 2019. The aorta was dissected and the aortic sinuses were opened, the number, location, size, shape and anomalous presence of accessory coronary ostia was noted and the measurements were taken using vernier calipers.

**Results:** The Right Coronary Artery (RCA) was seen taking origin from the anterior aortic sinus and the Left Coronary Artery (LCA) from the left posterior aortic sinus in all the 110 hearts. The mean diameter of the Left Coronary Ostia (LCO) was  $3.66 \pm 0.40$  mm and found to be greater than the Right Coronary Ostia (RCO)  $3.43 \pm 0.38$  mm. Variations in the location and shape of the RCO and LCO were noted, and presence of single accessory coronary ostia was observed in five cases. Presence of multiple accessory coronary ostia was observed in only one case and its presence was found to be very rare.

**Conclusion:** The diameter of the LCO was found to be greater than the RCO and the shape was circular in most of the coronary ostia. The majority of the coronary ostia were found to be located within the aortic sinuses below the Sinotubular Junction (STJ). Anatomical knowledge of the coronary ostia in the ascending aorta is very important during cardiac interventions around the aortic root.

**Keywords:** Aortic sinus, Osteological study, Third coronary artery

## INTRODUCTION

The origin of RCA and LCA of the heart is from the aortic sinus of Valsalva, the right arises from the anterior aortic sinus and the left from the left posterior aortic sinus, the right posterior aortic sinus is devoid of any orifice hence, known as a non-coronary sinus. The junction between the aortic dilatation and the aortic tube is known as sinotubular junction, this is represented by a well-defined circumferential supra-valvular ridge, and the coronary ostia normally open near to this ridge [1]. The morphometry of the coronary ostia is very important for various cardiovascular interventions such as cardiac catheterisation, percutaneous aortic root valve replacement, aortotomy, approach for aortic root enlargement, and surgical ostioplasty [2]. With the recent change in the lifestyle and increased stress among the various populations, coronary heart diseases are on the high rise, and several angiographies and coronary by-pass surgeries are being done on day to day basis in every hospital. According to the World Health Organisation (WHO), CHD constitutes the major cause of death in the industrial world, the risk factors being hypertension, diabetes, hyperlipidemia, and lack of physical exercise and other factors that can lead to the destruction of the vascular cells. The study of coronary ostia helps design coronary catheters or cannulas [3].

The aim of this study was to describe the location, size, and shape of the coronary ostia in cadaveric hearts and to compare the same on both RCA and LCA. This can shed more light on the anatomical knowledge on the coronary ostia necessary for the therapeutic and diagnostic cardiac interventions by both cardiologists and radiologists.

## MATERIALS AND METHODS

This was a descriptive cross-sectional study. The study was conducted at the dissection hall, Department of Anatomy, JSS Medical College, Mysuru, Karnataka, India. The heart specimens

were obtained from the mortuary of Mysore Medical College and JSS Medical College, Mysuru, Karnataka, India. A total of 110 adult cadaveric normal human hearts were collected 90 males (82%) and 20 (18%) females with an average mean age of males 67 years and females 63 years. The adult cadaveric hearts were obtained from the Department of Forensic Medicine of JSS Medical College and Mysore Medical College and Research Institute, Mysuru, Karnataka state, during March 2017 to March 2019. A sample size of 110 was calculated using the formula  $4pq/l^2$ . The number of cadaveric hearts was collected depending on the availability of the cadavers at the time of postmortem in the mortuary.

**Ethical consideration:** The study protocol was reviewed and approved by the JSS Medical College Institutional Ethical Review Board. Permission to conduct the study was obtained by the Institutional Ethical Committee (JSS Institutional Ethical approval number: JSSMC/PG/6399/2016-17) of JSS Medical College, JSSAHER, Mysuru, on 01/02/2017.

**Inclusion criteria:** Human adult cadaveric hearts were utilised with no history of cardiovascular diseases or cardiac surgery/injury with intact coronary arteries and aortic root.

**Exclusion criteria:** Cadaveric hearts with a history of cardiac injuries/trauma, cardiac surgeries, stent insertions, and cardiomegaly were not utilised for this study.

**Data entry:** All the morphological and morphometric measurements like the location, size and shape of both the coronary ostia were entered into the excel sheet by the investigator. The data was verified by rechecking all the parameters and validity ensured by randomly selecting 10% of observations and an average of three means were taken after each measurement of a parameter; this was cross-checked again while entering the data for any errors.

**Procedure:** The thoracic cavity was exposed after cutting the ribs and sternum, the parietal pericardium was incised along with the

great vessels to remove the heart from the thoracic cavity. Each specimen was immediately washed thoroughly in running tap water to remove all the blood clots and debris from the major vessels of the heart. The aorta and pulmonary trunk were identified and incised transversely about 3-4 inches above the supra-valvular ridge. The aortic root was incised vertically and the three aortic sinuses were identified along with their coronary ostia. A vertical incision was made along the tubular wall of the non-coronary sinus for clear visualisation of both the ostia. A vernier caliper was used to measure the diameters of RCO and LCO. The presence of any single or multiple coronary ostia was observed. In the present study, the following observations were recorded and analysed statistically.

1. The diameter of the coronary ostia;
2. The location of coronary ostia;
3. Variations in the shape of the coronary ostia and
4. Presence of anomalous accessory coronary orifices.

### STATISTICAL ANALYSIS

All data was analysed using SPSS software, version 22.0. Descriptive statistics were obtained. Independent t-tests were done for continuous variables. Pearson's chi-square tests were done to study the coronary ostia and morphometric differences between RCA and LCA. The p-value less than 0.05 considered significant.

### RESULTS

The aortic valves did not show any variations; in all the 110 specimens the three aortic cusps were normal. No openings were observed in the non-coronary sinus.

**1. Diameter of coronary ostia:** The average diameter of the RCO was 3.43±0.389 mm and the LCO was 3.66±0.40 mm. [Table/Fig-1] shows the measurement of the diameter of the coronary ostia being taken using the digital vernier calipers.



[Table/Fig-1]: Measurement of the coronary ostium using digital vernier calipers.


[Table/Fig-2] shows the comparison of the diameters of RCO and LCO. There is a statistically significant difference in diameters of RCO and LCO. The diameter of LCO was found to be greater than the RCO.

Side of coronary ostia	Frequency (n)	Mean (mm)	Standard deviation	p-value
Right	110	3.43	0.38	<0.001
Left	110	3.66	0.40	

[Table/Fig-2]: Comparison of the diameters of right and left coronary ostia.

\*Independent Sample t-test; n: Number of specimens

**2. Location of coronary ostia:** Normally the RCA arises from the right anterior aortic sinus and the LCA from the left posterior aortic sinus. In present study, the RCA in all the 110 hearts was seen taking origin from the anterior aortic sinus and the left from the left posterior aortic sinus. No anomalous origin from the non-coronary sinus was observed [Table/Fig-3].



[Table/Fig-3]: The normal location and shapes of coronary ostia.

STJ: Sinutubular junction; RAS: Right aortic sinus; NCS: Non-coronary sinus; LAS: Left aortic sinus

[Table/Fig-4] shows the frequency and percentage of variations in the location of RCO and LCO. The percentage of the presence of RCO and LCO within the sinus was found to be greater than the ostia present in the STJ. No openings were observed at the tubular level.


Side	Location of coronary ostia	Frequency (%)
Right	Sinus	85 (77.3)
	Sinutubular junction	25 (22.7)
	Tubular	0 (0)
	Total	110 (100)
Left	Sinus	81 (73.6)
	Sinutubular junction	29 (26.4)
	Tubular	0 (0)
	Total	110 (100)

[Table/Fig-4]: Showing the frequency and percentage of different locations of right and left coronary ostia.

**3. Variations in the shape of coronary ostia:** Three types of shapes have been observed like, Circular (C), Horizontally Elliptical (HE) and Vertically Elliptical (VE).

The [Table/Fig-5] heart specimen below shows the presence of VE right coronary ostium in the right aortic sinus.

The [Table/Fig-6] heart specimen below shows different shapes of coronary ostia at STJ, the RCO showing the circular shape (C) and the LCO showing HE shape of coronary ostia within the aortic sinus.



[Table/Fig-5]: The heart specimen showing the Vertically Elliptical (VE) coronary orifice. [Table/Fig-6]: The heart specimen showing the circular (C) and Horizontally Elliptical (HE) coronary Ostia seen at the Sinu-Tubular Junction (STJ). (Images from left to right)


The [Table/Fig-7] shows the frequency and percentage of variations in the shape of RCO and LCO. In 70.9% of cases the RCO was circular in shape, whereas the LCO was circular in 52.7% of cases, the highest frequency was seen with the circular shape, next was the HE and lastly the VE shape.

**4. Presence of a single accessory coronary ostia/Third Coronary Artery (TCA):** Single right accessory coronary ostia was present in 9% (10/110) and the LCO did not show any accessory ostia 0%. Double TCA was seen in three hearts (2.7%). [Table/Fig-8] shows the heart specimen with the anterior aortic sinus, with a single RCA Ostium and a single accessory coronary Ostium/TCA opening in the anterior aortic sinus in close relation to the RCA orifice.

Side	Shape of coronary ostia	Frequency (%)
Right	Horizontally elliptical	30 (27.3)
	Circular	78 (70.9)
	Vertically elliptical	02 (1.8)
	Total	110 (100)
Left	Horizontally elliptical	45 (40.9)
	Circular	58 (52.7)
	Vertically elliptical	07 (6.4)
	Total	110 (100)

**[Table/Fig-7]:** The frequency and percentage of different shapes of coronary ostia in present study.


**5. Presence of multiple small accessory coronary ostia:** Multiple right accessory ostia were seen in 7.27% (08/110) and there were no multiple ostia in the left coronary sinus 0%. The Sinoatrial (SA) nodal artery; vasa vasorum to the pulmonary trunk and ventricular branch of RCA was seen arising from the anterior aortic sinus apart from the TCA. Such a finding is rare and presence of this anastomotic arterial circle is an advantage since it indicates an important communication between the RCA and LCA. Totally, the presence of TCA was observed in 11.8% (13/110) of cases, 12 were male cadavers and one was female cadaver. There was no statistically significant difference in the parameters of the location, size, or shape of the coronary ostia among the males and females. [Table/Fig-9] shows the heart specimen showing the presence of RCA Ostia and accessory coronary orifice, presence of double accessory coronary ostia and presence of multiple accessory coronary ostia in the anterior aortic sinus. The probe in the figure represents the RCA Ostia.



**[Table/Fig-8]:** Showing the heart specimen with a single accessory coronary ostia (1) in close relation to the right coronary orifice (2).

**[Table/Fig-9]:** The heart specimens showing multiple accessory coronary ostia arising from the anterior aortic sinus along with the right coronary ostia. a: Single 3<sup>rd</sup> coronary artery; b: Right coronary ostia; c: Double 3<sup>rd</sup> coronary artery; d: Right coronary ostia; e: Multiple accessory coronary ostia. (Images from left to right)

[Table/Fig-10] depicts the percentage of the multiple accessory coronary ostia arising from the anterior aortic sinus along with the TCA in present study. The incidence of TCA was 11.8% and other orifices for vasa vasorum to the pulmonary trunk, SA nodal artery and ventricular branch of RCA were found to be 2.7% in present study. In all the 110 hearts, the accessory orifices were observed in the anterior aortic sinus in close relation to the right coronary orifice.



**[Table/Fig-10]:** Graphical representation of multiple accessory ostia seen in anterior aortic sinus.

## DISCUSSION

The cardiac catheterisation of the coronary arteries is done for either therapeutic or diagnostic intervention. It is done for evaluation and treatment of various heart diseases like coronary or myocardial diseases, measurement of haemodynamics in the heart, measurement of saturation and cardiac output, evaluation of left ventricular function, evaluation and treatment of cardiac arrhythmias. It was Claude Bernard who first placed the catheters into the arteries and veins of the horses. In 1929, it was Forssmann and Cournand, the Nobel Laureates who performed the revolutionary catheterisation of human hearts. In 1959, it was Sones who introduced the selective catheterisation technique for coronary angiography. In 1967 Ampaltz described the femoral route of introducing selective coronary catheters through the perpendicularly placed coronary Ostia [4-6]. Judkin's or Amplatz cannulas are most commonly used for coronary arteriography in the recent years. The emergence of the coronary artery orifices is a result of more of ingrowth rather than an outgrowth of the arterial channels. The mechanism considered reinforcing the theory of ingrowth to the development of coronary arteries is:

- The role of neural crest cells;
- Growth factors: Vascular Endothelial Growth Factor (VEGF) family members [7].

A study conducted in 2017 by Surekha WM et al., mentions the following clinical significance of the anatomical knowledge of coronary ostia [8]. It has been found that the presence of anomalous ostia can cause difficulty in cannulation of vessels during open-heart surgeries; in coronary arteriography and angiography an aortic root injection of dye is given to locate many ostia, anomalous ostia can cause fatal outcome; multiple coronary ostia may be associated with cardiac abnormalities like hypertrophic cardiomyopathy; knowledge of accessory ostia is important to interpret angiographic findings correctly; ectopic origin of right conus artery from the right aortic sinus is also known as TCA, it forms a separate path of anastomosis between the RCA and LCA [8].

According to Prajapati BSK et al., the coronary ostia above the STJ remains open during systole and diastole, this causes continuous blood flow and decreases coronary insufficiency [9]. In contrast, Joshi SD et al., stated that the aortic cusp does not flatten against the walls of the aortic sinus even at maximum systolic pressure [10]. Therefore, the ostia above the STJ is not advantageous. A high take-off of the coronary artery can lead to serious clinical implications like myocardial infarction and sudden cardiac death. However, it has been suggested that the location of ostia within the aortic sinus or below the STJ is functionally advantageous [10].

In present study, also the LCO (3.66±0.40 mm) had a greater diameter than the RCO (3.43±0.38 mm). [Table/Fig-11] shows the comparison of the mean diameters of the coronary ostia of present study with the previous studies [2-4,11,12].

Studies (Publication year)	Population	Sample size	RCO (mm)	LCO (mm)
Cavalcanti JS et al., [11] (2003)	Brazil	51	3.46±0.94	4.75±0.93
Kaur D et al., [12] (2012)	North India	77	3.9±1.0	4.6±1.0
Kulkarni JP and Vaishali P [2] (2015)	North India	90	2.5±1.0	2.8±1.0
Jyothi SR and Dakshayani KR [3] (2017)	India	49	3.4±0.9	4.3±0.7
Iman S et al., [4] (2017)	East India	40	4.19±0.36	4.41±0.4
Present study (2021)	South India	110	3.43±0.38	3.66±0.40

**[Table/Fig-11]:** Comparison of the diameters of the coronary ostia among different population with present study [2-4,11,12].

In present study, the percentage of circular shape was found to be highest in both RCO (70.9%) and LCO (52.7%) compared to the other shapes. [Table/Fig-12] shows comparison of the percentages of different shapes of coronary ostia among various population [2,13,14].

Studies (Publication year)	Sample size	Population	Right coronary ostia (%)			Left coronary ostia (%)		
			HE	C	VE	HE	C	VE
Gosva F et al., [13] (2010)	100	Turkey	-	60	-	-	55	-
Kulkarni JP and Vaishali P [2] (2015)	90	North India	76	16	7	73	23	10
Jeshika SL et al., [14] (2019)	50	South Africa	24	52	24	60	30	10
Present study (2021)	110	South India	27.3	70.9	1.8	40.9	52.7	6.4

**[Table/Fig-12]:** Comparison of the percentages of different shapes of coronary ostia among various population with present study [2,13,14]. HE: Horizontally elliptical; C: Circular; VE: Vertically elliptical

A study conducted by Cihan A et al., observed the angiographic incidence of anomalous aortic origins of the coronary arteries in 44.7% of patients. This study suggested that such anomalous coronary ostia can help to design appropriate catheters for angiographies [15].

In present study also, all the accessory coronary ostia were situated below the supra-valvular ridge/STJ. Banchi A in 1904 reported the highest percentage being 71% of RCO located at STJ [16] and 48% of LCO located at STJ compared to the other locations [17]. Wilkins CE in 1988 observed only the tubular locations of both the RCO and LCO [18]. Reig J and Petit M reported 56% of RCO and LCO being located in the sinus region and at STJ, no tubular location were found [19]. In present study also no tubular locations were found in all 110 hearts. Bharambe VK and Arole V reported the STJ being the commonest position compared to other locations, RCO in 62% of cases and LCO in 66% of cases were seen located at STJ [20].

Schlesinger MJ et al., (1949) was the first author to describe the TCA in detail, he mentioned that it supplied the right ventricle and inter-ventricular septum [21].

In present study, the RCO (77.3%) and LCO (73.6%) were located within the sinus, the RCO (22.7%) and LCO (26.4%) were located at STJ and no tubular locations were observed. The [Table/Fig-13] below shows the comparison of percentages of different locations of coronary ostia of previous authors with present study [3,16,18-20]. [Table/Fig-14] shows comparison of the incidence of third coronary artery with other studies [3,17,19,20,22,23]. [Table/Fig-15] shows comparison of the anomalous origin of different arteries from the anterior aortic sinus [3,20,24].

Authors (Publication year)	RCO/LCO	Sinus (%)	STJ (%)	Tubular (%)
Banchi A [16] (1904)	RCO	10	71	19
	LCO	18	48	34
Wilkins CE et al., [18] (1988)	RCO	-	-	8
	LCO	-	-	8
Reig J and Petit M [19] (2004)	RCO	56	56	-
	LCO	56	56	-
Bharambe VK and Arole V [20] (2012)	RCO	22	62	16
	LCO	10	66	24
Jyothi SR and Dakshayani KR [3] (2017)	RCO	89.8	8.2	3
	LCO	38.7	53.1	8.2
Present study (2021)	RCO	77.3	22.7	0
	LCO	73.6	26.4	0

**[Table/Fig-13]:** Comparison of the percentages of different locations of the right and left coronary ostia of previous studies with present study [3,16,18-20].

Muriango M et al., (1997) reported the 3<sup>rd</sup> coronary ostia in the anterior aortic sinus in five hearts and a single coronary ostia in five hearts [25].

Author (Publication year)	Percentage (%)
Reig J and Petit M [19] (2004)	33-51
Kalpna R [17] (2004)	24
Stankovic I and Jesic M [23] (2004)	34.8
Koenig PR and Hijazi ZM [22] (2005)	50
Bharambe VK and Arole V [20] (2012)	22
Jyothi SR et al., [3] (2017)	10.2
Present study (2021)	11.8

**[Table/Fig-14]:** Comparison of the incidence of 3<sup>rd</sup> coronary artery with the previous studies [3,17,19,20,22,23].

Previous studies (Publication year)	Anomalous origin from the anterior aortic sinus
Mc Alpine WA [24] (1975)	Ventricular branch of RCA
Bharambe VK and Arole V [20] (2012)	SA nodal artery, circumflex branch, vasa vasorum to the pulmonary trunk, ventricular branch of RCA.
Jyothi SR and Dakshayani KR [3] (2017)	Ventricular branch of RCA.
Present study (2021)	SA nodal artery, vasa vasorum to the pulmonary trunk, ventricular branch of RCA.

**[Table/Fig-15]:** Comparison of the anomalous origin of different arteries from the anterior aortic sinus (apart from the 3<sup>rd</sup> coronary artery) [3,20,24].

In present study, it was observed that the SA nodal artery; vasa vasorum to the pulmonary trunk and ventricular branch of RCA arising from the anterior aortic sinus apart from the TCA. Such a finding is rare and presence of this anastomotic arterial circle (circle of Vieussens) is an advantage since it indicates the communication between the RCA and LCA. Manisha RD et al., in her study stated the incidence of TCA in 48 out of 150 hearts (32%), in 42 hearts (28%) with a single TCA, and in 6 hearts (4%) double TCA was noted. All were closely related to the RCO. In two cases, 4 ostia were observed, one for RCA, 2 for TCA and 1 for the pulmonary trunk [26]. Gajbe UL et al., conducted study in 30 cadaveric hearts and found the presence of TCA in 2 hearts (6%) and presence of anomalous origin of RCA, SA nodal artery and vasa vasorum of pulmonary trunk as 3 separate openings in the anterior aortic sinus in 3 hearts (10%) [27]. Urbanski PP et al., performed the ascending aorta replacement in patient with coronary ostia localised above the sinotubular junction by slide-plasty of an aberrant coronary artery. Transposition of the anomalous coronary artery origin from the replaced ascending aorta into the normal sinus of valsalva using slide-plasty was considered as a simple and safe surgical procedure for which a sound anatomical knowledge of the size, location and shape of the coronary Ostia is essential for the operating surgeon in the same population [28].

**Limitation(s)**

The major limitation was that the number of female cadaveric hearts was comparatively less than the male hearts. This study will help in interventional studies by cardiologist and radiologists during diagnostic and therapeutic approach in coronary vascular disease and its management. The present study of size, location and shape of the coronary ostia will also help to design appropriate catheters and cannulas for coronary angiography and cardiac interventions for the same population.

**CONCLUSION(S)**

The diameter of the LCO was found to be greater than the RCO and the shape was circular in most of the coronary ostia. The majority of the coronary ostia were found to be located within the aortic sinuses below the STJ. This anatomical study will enhance the clinical knowledge of the variations associated with the coronary ostia and will greatly help the radiologists performing coronary angiographies.

## Acknowledgement

Author would like to thank JSS Academy of Higher Education and Research, Mysuru, Karnataka state, India, for allowing to publish the research work. Author thank Head of Department Dr. Pushpalatha K, Department of Anatomy, JSS Medical College, JSS Academy of Higher Education and Research, for her support. Author is very grateful to guide and Principal Dr. H. Basavanagowdappa, JSS Medical College, JSS Academy of Higher Education and research, for his constant support and encouragement during my research work. Author is thankful to Dr. Madhu, Associate Professor, Community Medicine Department, JSS Medical College, JSS Academy of Higher Education and Research, for her immense support and guidance during the research work.

All the authors sincerely thank those who donated their bodies to science so that anatomical research could be performed. Results from such research can potentially increase mankind's overall knowledge that can then improve patient care. Therefore, these donors and their families deserve our highest gratitude.

## REFERENCES

- [1] Standing S, Ellis H, et al. Gray's anatomy. In thorax, 39<sup>th</sup> edition. Elsevier Churchill Livingstone, Edinburg, London, New York, Oxford, Philadelphia, St.Louis, Sydney, Toronto 2005; 1014-1018.
- [2] Kulkarni JP, Vaishali P. Topography, morphology and morphometry of coronary ostia- A cadaveric study. Eur J Anat. 2015;19(2):165-70.
- [3] Jyothi SR, Dakshayani KR. Morphometric study of coronary ostia in human cadavers by dissection method. Indian J Clin Anat Physio. 2017;4(2):130-32.
- [4] Iman S, Ratnadeep P, Alipta B, Asis KG, Akash KB, Prasanta C. The debatable aspect of the coronary ostia and their off- shoots: An attempt to solve it, based on cadavers in eastern India. Indian J Basic Applied Med Res. 2017;7(1):109-19.
- [5] Bagchi, Asoke K. Hinduja Foundation Encyclopaedia of Nobel Laureates, 1901-1987, Delhi: Konark Publishers Pvt., Ltd.; 1988; pp 364.
- [6] Gensini GG. Historic events in the development of coronary arteriography. In: Coronary Arteriography. New York: Futura Publishing Company; 1975; pp 163.
- [7] Silva Junior GO, Miranda SWS, Mandarim de Lacerda C. Origin and development of coronary arteries. Int J Morphol. 2009;27(3):891-98.
- [8] Surekha WM, Anjali RL, Rukhmode VP, Ujwal LG. Multiple Coronary Ostia and their clinical and surgical significance. ISOR Journal of Dental and Medical Sciences. 2017;16(5):01-08.
- [9] Prajapati BSK, Patil D, Udajna A, Bhatt CPV. Variation in ostium of coronary arteries. Natl J Med Res. 2013;3:134-36.
- [10] Joshi SD, Joshi SS, Athavale SA. Origins of the coronary arteries and their significance. Clinics Journal. 2010;65:79-84.
- [11] Cavalcanti JS, De Melp NC, Vasconcelos RS. Morphometric and topographic study of coronary ostia. Arq Bras Cardiol. 2003;81(4):355-62.
- [12] Kaur D, Singh K, Nair N, Kalra. Morphology and morphometry of coronary ostia in South Indian adult human hearts. Int J Biol Med Res. 2012;3(3):2169-71.
- [13] Govsa F, Celik S, Aktas EO, Aktas S, Kocak A, Boydak B, et al. Anatomic variability of the coronary arterial orifices. Anatolian J Cardio. 2010;10(1):03-08.
- [14] Jeshika SL, Luckrajh, Kapil SS, Lelika L. An anatomical investigation of the coronary ostia and its relationship to the sinotubular junction within a select South African population. Eur J Anat. 2019;23(3):159-65.
- [15] Cihan A, Suleyman K, Koc S, Gursoy YC, Bal U, Aydinalp A, et al. Coronary anatomy, anatomic variations & anomalies: A retrospective coronary angiography study. Singapore Med J. 2015;56(6):339-45.
- [16] Banchi A. Morfologia della arteriae coronariae cordis. Arch Ital Anat Embryol. 1904;3:87-164.
- [17] Kalpana R. A study on principal branches of coronary arteries on humans. J Anat Soc India. 2005;52(2):137-40.
- [18] Wilkins CE, Benjamin B, Virendra SM, Ali M, Carlos MDC, Garcia E, et al. Coronary anomalies. Tex Heart Institute Journal. 1988;15:166-73.
- [19] Reig J, Petit M. Main trunk of the left coronary artery: Anatomical study of the parameters of clinical interest. Clin Anat. 2004;17:06-13.
- [20] Bharambe VK, Arole VA. A study of variations in coronary ostia. J Anat Soc India. 2012;61(2):221-28.
- [21] Schlesinger MJ, Zoll PM, Wessler S. The conus artery: A third coronary artery. Am Heart J. 1949;38:823-38.
- [22] Koenig PR, Hijazi ZM. Congenital coronary artery abnormalities. In uptodate (Edn Rose BD) Waltham, Mass: Uptodate, 2005.
- [23] Stankovic I, Jestic M. Morphometric characteristics of the conal coronary artery. MJM. 2004;8:02-06.
- [24] Mc Alpine WA. Heart and coronary arteries Berlin: Springer- Verlag; 1975. pp 133-209.
- [25] Muriago M, Sheppard MN, Ho SY, Anderson KH. Location of the coronary arterial orifices in the normal heart. Clin Anat. 1997;10(5):297-302.
- [26] Manisha RD, Medha GP, Uttama UJ. Study of third coronary artery in adult human cadaveric hearts. J Clin Diag Res. 2015;9(10):01-04.
- [27] Gajbe UL, Gosavi S, Meshram S, Gajbhiye VM. The anomalous origin of multiple coronary ostia and their clinical significance. J Clin Diag Res. 2010;(4):2129-33.
- [28] Urbanski PP, Irimie V, Diegeler A, Morka A, Thamm T, Lehmkuhl L, et al. Ascending aorta replacement in patients with coronary ostia localised above the sinotubular junction. European Journal of Cardio-Thoracic Surgery. 2020; <https://doi.org/10.1093/ejcts/ezaa.414>.

### PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of Anatomy, JSS Medical College, Mysuru, Karnataka, India.
2. Professor, Department of Medicine, JSS Medical College, Mysuru, Karnataka, India.
3. Associate Professor, Department of Community Medicine, JSS Medical College, Mysuru, Karnataka, India.

### NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Najma Mobin,  
#808, S.S. Nagar, Bannimantap, Mysuru-570015, Karnataka, India.  
E-mail: drnajmamobin786@gmail.com

### PLAGIARISM CHECKING METHODS: [\[Jain H et al.\]](#)

- Plagiarism X-checker: Nov 25, 2020
- Manual Googling: Dec 28, 2020
- iThenticate Software: Jan 05, 2021 (9%)

### ETYMOLOGY: Author Origin

### AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? No
- For any images presented appropriate consent has been obtained from the subjects. NA

Date of Submission: **Nov 22, 2020**

Date of Peer Review: **Dec 22, 2020**

Date of Acceptance: **Jan 05, 2021**

Date of Publishing: **Feb 01, 2021**