

Analysing the Functional and Structural Impact of Single Nucleotide Polymorphisms in Matrix Metalloproteinase 3 Gene: An In-silico Approach

REJITHA RAMACHANDRAN¹, SARITHA C JOSEPH², KC SAJEEVAN³, SHAJEE S NAIR⁴

ABSTRACT

Introduction: Matrix Metalloproteinase 3 (MMP3) is a vital member of the MMP family, known for its wide range of substrate specificity and proteolytic activity against Extracellular Matrix (ECM) components. The role of functional polymorphisms in the MMP genes has been previously investigated in relation to cancer susceptibility, particularly breast cancer. Several Single-Nucleotide Polymorphisms (SNPs) in the MMP3 gene have been linked to a number of clinical illnesses, such as Coronary Artery Disease (CAD); however, the results were not entirely conclusive.

Aim: To identify pathogenic missense SNPs in the human MMP3 gene and analyse their effects on structure and function.

Materials and Methods: This was a record-based cross-sectional study performed using data retrieved from online resources. The analysis was conducted using a series of different bioinformatic tools, for which ethical clearance was obtained from the institution. The online tools used included Sorting Intolerant from Tolerant (SIFT), PolyPhen-2, PhD-SNP, PANTHER, PROVEAN, and SNPs

and GO to predict harmful non synonymous SNPs (nsSNPs). Further analysis was performed using I-Mutant 2.0, MutPred2, ConSurf, and HOPE software. These tools were able to filter out damaging SNPs and predict the impact of deleterious SNPs on the structure and function of the MMP3 protein.

Results: This study predicted two potentially pathogenic SNPs (D175Y and Y116C) out of 443 missense SNPs from dbSNP, which is a database of SNPs available on the National Centre for Biotechnology Information website. Further analysis revealed that these SNPs were located in highly conserved regions and were predicted to decrease protein stability.

Conclusion: In this study, two potentially pathogenic SNPs (D175Y and Y116C) were identified. Characterisation of these SNPs can help us gain a better understanding of the molecular basis of clinical conditions. The results of this study can be further validated by designing population-based studies and wet lab experiments. This will help in augmenting research and personalised medicine.

Keywords: Computational molecular biology, Gene polymorphism, Human genome, Precision medicine

INTRODUCTION

The human genome is relatively large, comprising approximately six billion nucleotides. About 99.9% of the DNA sequences in the human genome are the same across individuals, with individual variations accounting for the remaining 0.1% [1]. Single Nucleotide Polymorphisms (SNPs), which refer to the substitution of a single nucleotide, are the most prevalent type of genetic variation. SNPs occur throughout the genome at a rate of one in every 1,000 base pairs. SNPs are broadly categorised into two types: synonymous and non synonymous (nsSNP). Non synonymous SNPs are responsible for protein variations and amino acid substitutions in humans, while synonymous SNPs do not change the amino acid sequences [2]. With the development of practical and readily accessible DNA sequencing tools, SNPs have gained substantial clinical significance over the past two decades. Regardless of their effects on the biological function of gene products, SNPs are increasingly recognised as valuable markers in mapping complex human diseases, population genetics, and evolutionary studies [3]. Efficient identification of important SNPs and understanding their impact on protein function will be beneficial for genetic studies aimed at elucidating the molecular basis of various diseases. Furthermore, targeted in-vitro and in-vivo experiments can be designed to predict the effects of specific SNPs [4]. In light of this, multiple studies are being conducted to predict the functional consequences of nsSNPs, particularly based on sequence information and structural attributes [5-10].

Matrix Metalloproteinases (MMPs) are a family of more than 25 zinc-containing enzymes that break down various proteoglycans,

fibronectin, interstitial collagen, and basement membrane collagen during normal physiological remodelling and repair processes in development and inflammation [11]. MMP-3, also known as Stromelysin-1, is a crucial member of the MMP family that exhibits a wide range of substrate specificity. Additionally, MMP-3 plays a critical role in the remodelling of connective tissue by activating other MMPs, such as collagenase, matrilysin, and gelatinase B [12,13]. The role of MMP gene functional polymorphisms in cancer susceptibility, especially breast cancer, has been previously investigated. The 5A/6A polymorphism has been linked to patient survival, oestrogen receptor status, breast cancer risk, and tumour size based on multiple genome-wide association analyses [14]. However, the association between SNPs and breast cancer risk remains inconclusive and controversial, as it is largely influenced by the ethnicity of the population and sample size [14-16]. Apart from breast cancer, a meta-analysis indicates a significant correlation between gastrointestinal, colorectal, and oesophageal malignancies and the MMP-3-1171 5A/6A variation [17].

In addition to malignancies, MMP3 gene polymorphisms have been found to be associated with atherosclerosis. This is due to the critical role of MMP3 in the degradation of the Extracellular Matrix (ECM) of blood vessels. Several studies evaluating the influence of promoter polymorphism and MMP-3 (-1612 5A/6A) on CAD risk have produced contradictory results, including gender differences as well [18-20]. Two MMP3 polymorphisms, rs520540 A/G and rs679620 C/T, were associated with cerebral stroke risk; however, contrary results were obtained in a different population, probably due to racial differences [21].

Among these reported SNPs, distinguishing functional SNPs from neutral SNPs is a major challenge. Experimental wet lab studies aimed at identifying the effects of multiple SNPs are time-consuming, challenging, and costly. Applying computational approaches to filter out possibly detrimental SNPs is emerging as a novel strategy for understanding the molecular mechanisms of diseases [22].


This was an in-silico study utilising bioinformatics computational tools with good predictive capabilities to identify the possible deleterious SNPs in the MMP-3 gene. Several studies have been conducted to find the association between specific SNPs in the MMP3 gene and diseases [14-21]. However, according to the literature search, there have been no comparable in-silico studies that examined every SNP in MMP-3 from the NCBI database. This study is expected to unravel the potential impact of polymorphisms on MMP3 structure and function. In addition, it can help in designing further specific experiments to confirm the importance of these SNPs in various clinical conditions. Combined with experimental data, it can aid in the development of personalised medicine.

MATERIALS AND METHODS

This online records-based cross-sectional study was conducted at Government Medical College, Manjeri, Kerala, India over a period from May to July 2024 (three months), which included data collection as well as analysis using online computational tools. IEC approval for this study was obtained from the institution (Ref. No: IEC/GMCM/133, date: 18/05/2024).

Methodology

The operational flow chart of this study is presented in [Table/Fig-1].



S. No.	Tool	website	Based on	Input	Output
1	SIFT	https://sift.bii.a-star.edu.sg/	Sequence homology and the physical properties of amino acids	rsID	Score and prediction
2	Polyphen2	http://genetics.bwh.harvard.edu/pph2/	Structural and comparative evolutionary considerations	uniprotID or fasta protein sequence, amino acid change	Score and prediction
3	Phd-SNP	http://snps.biofold.org/phd-snp/phd-snp.html	Supervised learning based	FASTA sequence, swiss-prot code, amino acid change	Ri value and prediction
4	PANTHER	https://www.pantherdb.org/	Sequence homology	FASTA sequence and substitutions	PSEP score and message
5	PROVEAN	http://provean.jcvi.org/index.php	Sequence homology	rsID	Score and prediction
6	SNPs&GO	https://snps.biofold.org/snps-and-go/snps-and-go.html	Supervised learning based	FASTA sequence, swiss-prot code, amino acid change	Ri value and prediction
7	MUTPRED2	http://mutpred.mutdb.org	Machine learning-based method	Fasta sequence and amino acid change	Pathogenicity prediction, and a ranked list of specific molecular alterations
8	I-MUTANT	https://folding.biofold.org/i-mutant/	Support vector machine -based web server	Protein sequence and amino acid change	DDG value and reliability index

The detailed methodology is as follows:

- 1. Retrieval of SNPs in the MMP3 gene:** The ncbi SNPs database (dbSNP) (<https://www.ncbi.nlm.nih.gov/snp/>) was used to retrieve all known SNPs for MMP-3 in humans [23]. Non synonymous (missense) mutations were selected from the dataset for further analysis. The UniProt database was utilised to obtain the protein's sequence and other pertinent information.
- 2. Prediction of the most deleterious missense SNPs:** The retrieved missense SNPs were analysed using bioinformatics-based web tools for identification of high-risk nsSNPs. Details of the tools used are provided in [Table/Fig-2]. Polymorphism phenotyping v2 (PolyPhen-2) and SIFT were the initial web-based tools employed. A SIFT score value between 0 and 1 indicates a potentially harmful effect, with scores below 0.05 being particularly concerning [24]. A PolyPhen score ranging from 0.85 to 1 is considered probably damaging [25,26]. SNPs identified as potentially harmful by the aforementioned tools were further subjected to additional bioinformatics tools for confirmation and further refinement [27]. The tools used included Predictions from Predictor of Human Deleterious Single Nucleotide Polymorphisms (PhD-SNP) [28], PANTHER-position-specific evolutionary preservation [29], PROVEAN [30], and SNPs and GO [31]. SNPs projected by every tool to be harmful were selected for further examination.
- 3. Structural and functional analysis of SNPs with predicted damaging consequences:**
 - a. Identification of impact on the structural and functional properties of MMP3:** MutPred2 is a web application that integrates genetic and molecular data. It provides a ranked list of specific molecular changes with potential impacts on phenotype, along with a general pathogenicity prediction [32].
 - b. Prediction of change in stability of the MMP3 protein using I-Mutant 2.0 Suite:** The influence of nsSNPs on protein stability was assessed using the I-Mutant 2.0, which utilises the free energy change value (Delta Delta G: DDG). The DDG represents the difference between the free energy of the folded wild-type and mutant structures. DDG values fall into three categories: neutral ($-0.5 \leq DDG \leq 0.5$ kcal/mol), mostly stable ($DDG > 0.5$ kcal/mol), or highly unstable ($DDG < -0.5$ kcal/mol) [33].
 - c. Prediction of evolutionary conservation of MMP3 using the ConSurf server:** The ConSurf web server was used to examine the evolutionary rate of conservation of amino acids in MMP3. ConSurf determines the conservation scores of the protein's amino acid positions by analysing the evolutionary relationships among similar

9	CONSURF	https://consurf.tau.ac.il	Reconstruction of phylogenetic tree reflecting evolutionary relations	Uniprot ID	Amino acid sequence conservation scores
10	HOPE	https://www3.cmbi.umcn.nl/hope	BLAST, Homology modelling, information from Uniprot and Refseq	Unitprot ID, Fasta sequence and amino acid change	Effect of SNP on protein structure
11.	MUTATION3D	http://www.mutation3d.org/	Complete-linkage clustering	unitprotID, amino acid change	3D structure and linear protein models

[Table/Fig-2]: Details of the study's bioinformatics tools.

sequences. The highest conserved location is graded as 9, while conservation values range from 1 to 9 [34].

- d. **Protein structure prediction and mutant analysis using HOPE software:** The web-based application Have (y) Our Protein Explained (HOPE) is used to analyse how SNPs affect the structural conformation and functionality of proteins. HOPE uses data from multiple web services and databases to generate reports regarding the impact of SNPs, which include text, graphs, and animations [35].

RESULTS

1. **Data on SNPs retrieved:** In the MMP3 gene, the dbSNP yielded a total of 4,120 SNPs. Among these SNPs, 443 were missense, 196 were synonymous, and 2,720 were located in the intronic region. For further in-silico research, only the missense or nsSNPs were selected. The remaining 761 SNPs belonged to other functional classes, for which authors were unable to retrieve information from dbSNP. The functional classes in dbSNP

SI no	SNP rsID	Amino acid change	SIFT score	Polyphen-2 score
1	rs185081958	A391V	0.012	0.999
2	rs373335227	Y401H	0.001	0.992
3	rs201409153	Y344C	0.008	0.959
4	rs375381761	L294W	0	0.997
5	rs147533686	R316C	0	1
6	rs145941285	I159T	0.001	0.976
7	rs150552863	I220T	0.004	0.938
8	rs201427128	D175Y	0.015	0.92
9	rs140794544	P187S	0	0.993
10	rs141650167	R91W	0	1
11	rs61753776	W109G	0	0.996
12	rs142984123	Y116C	0.01	1
13	rs373783506	V144M	0.001	0.994
14	rs374032529	L10P	0.003	0.919

[Table/Fig-3]: Results of SIFT and Polyphen2 bioinformatics tools.

Amino acid change	Phd-SNP		PROVEAN		PANTHER			SNPs and GO	
	Prediction	RI	Score	Prediction	Preservation time	Message	Pdel	Prediction	RI
A391V	Neutral	7	-3.661	Deleterious	911	Probably damaging	0.85	Neutral	6
Y401H	Disease	4	-4.248	Deleterious	361	Possibly damaging	0.5	Neutral	6
Y344C	Disease	5	-7.689	Deleterious	455	Probably damaging	0.57	Neutral	6
L294W	Neutral	7	-5.189	Deleterious	456	Probably damaging	0.57	Neutral	5
R316C	Neutral	1	-7.293	Deleterious	797	Probably damaging	0.74	Neutral	1
I159T	Disease	1	-4.934	Deleterious	911	Probably damaging	0.85	Neutral	4
I220T	Disease	5	-2.424	Neutral	176	Probably benign	0.27	Neutral	8
D175Y	Disease	5	-8.686	Deleterious	911	Probably damaging	0.85	Disease	3
P187S	Disease	2	-7.858	Deleterious	911	Probably damaging	0.85	Neutral	2
R91W	Neutral	1	-7.527	Deleterious	911	Probably damaging	0.85	Disease	3
W109G	Neutral	1	-12.841	Deleterious	911	Probably damaging	0.85	Disease	3
Y116C	Disease	3	-8.879	Deleterious	911	Probably damaging	0.85	Disease	2
V144M	Disease	2	-2.593	Deleterious	797	Probably damaging	0.74	Neutral	5
L10P	Disease	3	-4.604	Deleterious	176	Probably benign	0.27	Neutral	3

[Table/Fig-4]: Results of SNPs chosen to be damaging by SIFT and Polyphen when subjected to other bioinformatics tools.

delineate the position of a polymorphism in relation to identifiable features of a specific gene transcript. Most functional classes are determined by the location of the variation in relation to the exon boundaries of the transcript [23]. Since present study requires only missense SNPs, only those were retrieved and analysed. The remaining 3,677 SNPs were excluded from the manuscript due to our inability to obtain accurate information about their functional class.


2. **Identification of predicted deleterious SNPs:** The selected missense SNPs were subjected to SIFT and PolyPhen-2 to identify possible deleterious SNPs. Out of the 443 SNPs, 14 were predicted to be damaging by both SIFT and PolyPhen-2 [Table/Fig-3] [24-26]. All 14 shortlisted SNPs identified by these two web-based techniques were further analysed using other bioinformatics tools: Phd-SNP, PANTHER, PROVEAN, and SNPs and GO [Table/Fig-4] [28-31]. Among all the analysed SNPs, two nsSNPs, rs201427128 (D175Y) and rs373783506 (Y116C), were predicted to be damaging by all six bioinformatics tools, including SIFT and PolyPhen.
3. **Prediction by MutPred2 on structural and functional effects [32]:** The selected two detrimental SNPs were analysed using MutPred2, which suggested a significant impact on the 3D structure of the MMP3 protein [Table/Fig-5].
4. **Prediction of changes in protein stability by I-Mutant 2.0 [33]:** I-Mutant predicted the extent to which the damaging SNPs alter the stability of the MMP3 protein. Both SNPs were predicted to decrease the stability of the protein [Table/Fig-6].
5. **Consurf server results [34]:** An analysis of conservation using the ConSurf server revealed that both SNPs, D175Y and Y116C, were located in highly conserved areas and were found to be exposed residues [Table/Fig-7].
6. **Predicted changes in the 3D structure of the MMP3 protein due to nsSNPs [35]:** HOPE software was used to generate results for both nsSNPs, with structural information obtained from the Protein Data Bank (PDB) database. The PDB ID used was 1SLM. In the case of D175Y, the mutant residue is larger, impacting hydrogen bond formation and its

SNP	Molecular changes (p-values <=0.05)	MutPred score
D175Y	Altered metal binding, ordered interface and Transmembrane protein Loss of Catalytic site at D175 Loss of Relative solvent accessibility Loss of Allosteric site at F174 Loss of Sulfation at Y172	0.921
Y116C	Alteration in ordered interface, Metal binding, DNA binding, Transmembrane protein Loss of Allosteric site at Y121 Altered stability	0.849

[Table/Fig-5]: Results from MutPred analysis.


S. No.	SNP	DDG (Kcal/mol)	RI
1	rs201427128 (D175Y)	-1.40	3
2	rs373783506 (Y116C)	-0.03	2

[Table/Fig-6]: Results of I-mutant 2.0 analysis of nsSNPs.



[Table/Fig-7]: ConSurf results for MMP3 protein.

interaction with the metal ion in the original wild-type residue. Additionally, there is a disturbance in the ionic interaction made by the wild-type residue. For Y116C, the smaller residue results in an empty space in the protein core, which disrupts proper folding by causing a loss of hydrogen bonds in the core [Table/Fig-8].



[Table/Fig-8]: a) An overview of the protein is presented in a ribbon format, with colours indicating different structural elements: α -helix in blue, β -strand in red, turn in green, 3/10 helix in yellow, and random coil in cyan; b) A close-up view of the D175Y mutation; c) A close-up view of the Y116C mutation. The protein is rendered in grey, while the side chains of both the wild-type and mutant residues are highlighted in green and red, respectively.

DISCUSSION

The MMP-3 gene, which codes for the MMP-3 protein, is a component of the MMP gene cluster located on chromosome 11q22.3. The NCBI gene ID for MMP3 is 4314, and its UniProt ID is P08254 [36,37]. The expression of the MMP3 gene may be regulated by DNA polymorphisms, which can exert allele-specific effects, resulting in fundamental variations in each person's susceptibility to diseases and other traits [11].

The goal of the current study was to use several computational techniques to discover SNPs that could be harmful and alter the structure or function of the MMP3 gene. A significant number of researchers have utilised in-silico tools to predict the structural and functional impact of SNPs on proteins, such as the human CTLA4 gene [38], IL-33 gene [39], and the androgen receptor gene [27]. This work represents the first comprehensive and systematic in-silico study of functional SNPs in the MMP3 gene.

Combining multiple computational methods can enhance the validity of predictions regarding damaging SNPs and improve efficiency. By employing six prediction methods (SIFT, PolyPhen-2, PROVEAN, PANTHER, PhD-SNP, and SNPs and GO), authors were able to obtain an integrated view of harmful SNPs in the MMP3 gene. Since each method relies on a different set of parameters, two ns SNPs (D175Y and Y116C) were identified as high-risk in present study, which were predicted to be harmful by every algorithm. Additional research revealed that these two ns SNPs might be dangerous because they are located in a highly conserved region and have the potential to affect protein stability [6-9,22,27,38,39].

As an important member of the MMP3 family, several experimental studies have been conducted on genetic variation in MMP3, especially in malignancies [11,13-18,21]. Wang L et al., conducted an observational study and meta-analysis of MMP3 gene polymorphism in the Chinese population. The researchers concluded that while MMP-3 gene polymorphism raises the risk of ovarian cancer in the southern Chinese population, their meta-analysis showed that it had no effect on risk in other populations [40]. Durmanova V et al., demonstrated that the MMP3 SNP rs3025058 influences the age of onset of Alzheimer's disease [41].

The predicted ns SNPs from present study have not been linked to any clinical disease in published studies. Therefore, to support this finding, it is necessary to validate these ns SNPs through larger population-based studies and specifically designed wet lab experiments.

Limitation(s)

The present study utilised the limited dataset from the NCBI dbSNP database. More ns SNPs could have been analysed if other databases had been included.

CONCLUSION(S)

Using a variety of bioinformatics techniques, this study has identified two significant nsSNPs, D175Y and Y116C, from the extensive SNP dataset of MMP3. Based on these findings, it can be concluded that these SNPs should be regarded as significant candidates for planning functional experiments involving the MMP3 gene. This presents an opportunity to investigate the molecular basis of any associated clinical problems and may aid in the discovery of potential drugs or pharmacological targets. Although computational biology is powerful, it has its limitations; therefore, population-based research is necessary to provide definitive validation.

REFERENCES

- Teama S. DNA Polymorphisms: DNA-Based molecular markers and their application in medicine [Internet]. Genetic Diversity and Disease Susceptibility. InTech; 2018. Available from: <https://dx.doi.org/10.5772/intechopen.79517>.
- Hossain MS, Roy AS, Islam MS. In silico analysis predicting effects of deleterious SNPs of human RASSF5 gene on its structure and functions. Sci Rep. 2020;10(1):14542.

- [3] Wakui M. Analysis of Single Nucleotide Polymorphisms (SNPs). *Rinsho Byori*. 2013;61(11):1008-17.
- [4] Chitrana KN, Yeguvapalli S. Computational screening and molecular dynamic simulation of breast cancer associated deleterious non-synonymous single nucleotide polymorphisms in TP53 gene. *PLoS one*. 2014;9(8):e104242.
- [5] George Priya Doss C, Rajasekaran R, Sudandiradoss C, Ramanathan K, Purohit R, Sethumadhavan R. A novel computational and structural analysis of nsSNPs in CFTR gene. *Genomic Med*. 2008;2(1-2):23-32.
- [6] Nouria S, Bouatrous E, Ouragini H. An in-silico analysis of the coding and non-coding non-synonymous single nucleotide variants in the Npm1 gene to determine susceptibility to cancer. 2024. Available from: <https://doi.org/10.2139/ssrn.4871550>.
- [7] Almakhari M, Chen Y, Kong AS, Moradigaravand D, Lai KS, Lim SH, et al. In-silico identification of deleterious non-synonymous SNPs of TBX1 gene: Functional and structural impact towards 22q11. 2DS. *Plos one*. 2024;19(6):e0298092.
- [8] Subramanian S, Sanjeev K, Prakash PS, Appukkuttan D. In silico analysis of single nucleotide polymorphisms of NAT-2 gene and its implications in periodontitis. *Int J Chem Biochem Sci*. 2023;24(8):177-81.
- [9] Wiguna NY, Hawa P, Jonny J. Human PPARG gene mutation as risk factor for Type 2 diabetes mellitus: In silico analysis. *ASEAN Journal of Military and Preventive Medicine*. 2024;1(1):01-02.
- [10] Abdelazim AA, Maged M, Abdelmaksoud AI, Hassanin SE. In-silico analysis of nsSNPs in human CYP3A4/5 and their effects on drug-enzymes interactions of FDA-approved COVID-19 antiviral drugs: A pharmacogenetic study. 2024. Doi:10.21203/rs.3.rs-3910067/v1.
- [11] Munhoz, Godoy-Santos AL, Santos MC. MMP-3 polymorphism: Genetic marker in pathological processes. *Mol Med Rep*. 2010;3(5):735-40.
- [12] Cabral-Pacheco GA, Garza-Veloz I, Castruita-De la Rosa C, Ramirez-Acuña JM, Perez-Romero BA, Guerrero-Rodríguez JF, et al. The roles of matrix metalloproteinases and their inhibitors in human diseases. *Int J Mol Sci*. 2020;21(24):9739.
- [13] Ye S, Eriksson P, Hamsten A, Kurkinen M, Humphries SE, Henney AM. Progression of coronary atherosclerosis is associated with a common genetic variant of the human stromelysin-1 promoter which results in reduced gene expression. *J Biol Chem*. 1996;271(22):13055-60.
- [14] Suhaimi SA, Chan SC, Rosli R. Matrix Metalloproteinase 3 polymorphisms: emerging genetic markers in human breast cancer metastasis. *J Breast Cancer*. 2020;23(1):01-09.
- [15] Peng B, Cao L, Wang W, Xian L, Jiang D, Zhao J, et al. Polymorphisms in the promoter regions of matrix metalloproteinases 1 and 3 and cancer risk: A meta-analysis of 50 case-control studies. *Mutagenesis*. 2010;25(1):41-48.
- [16] Ghilardi G, Biondi ML, Caputo M, Leviti S, DeMonti M, Guagnellini E, et al. A single nucleotide polymorphism in the matrix metalloproteinase-3 promoter enhances breast cancer susceptibility. *Clin Cancer Res*. 2002;8(12):3820-23.
- [17] Aziz MA, Jafri S, Barek MA, Anonna SN, Islam MS. MMP-3 -1171 5A/6A promoter polymorphism and cancer susceptibility: An updated meta-analysis and trial sequential analysis. *Future Oncol*. 2023;19(21):1495-512.
- [18] Wenwang L. Susceptibility of MMP3 gene polymorphism to coronary artery disease: A meta-analysis. *J Med Biochem*. 2023;42(4):685-93.
- [19] Beyzade S, Zhang S, Wong YK, Day IN, Eriksson P, Ye S. Influences of matrix metalloproteinase-3 gene variation on extent of coronary atherosclerosis and risk of myocardial infarction. *J Am Coll Cardiol*. 2003;41(12):2130-37.
- [20] Koch W, de Waha A, Hoppmann P, Schömig A, Kastrati A. Haplotypes and 5A/6A polymorphism of the matrix metalloproteinase-3 gene in coronary disease: Case-control study and a meta-analysis. *Atherosclerosis*. 2010;208(1):171-76.
- [21] Yin Y, Zhang Y, Zhang X, Zhang Q, Wang J, Yang T, et al. Association of MMP3, MMP14, and MMP25 gene polymorphisms with cerebral stroke risk: A case-control study. *BMC Med Genomics*. 2023;16(1):297.
- [22] Desai M, Chauhan J. In silico analysis of nsSNPs in human methyl CpG binding protein 2. *Meta Gene*. 2016;10:01-07.
- [23] Bhagwat M. Searching NCBI's dbSNP database. *Curr Protoc Bioinformatics*. 2010;32(1):01-09.
- [24] Sim NL, Kumar P, Hu J, Henikoff S, Schneider G, Ng PC. SIFT web server: Predicting effects of amino acid substitutions on proteins. *Nucleic Acids Res*. 2012;40(W1):W452-57.
- [25] Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense mutations using PolyPhen-2. *Curr Protoc Hum Genet*. 2013;76(1):07-20.
- [26] Gemovic B, Perovic V, Glisic S, Veljkovic N. Feature-based classification of amino acid substitutions outside conserved functional protein domains. *Scientific World J*. 2013;2013(1):948617.
- [27] Adiba M, Das T, Paul A, Das A, Chakraborty S, Hosen MI, et al. In silico characterization of coding and non-coding SNPs of the androgen receptor gene. *Informatics in Medicine Unlocked*. 2021;24:100556.
- [28] Capriotti E, Calabrese R, Casadio R. Predicting the insurgence of human genetic diseases associated to single point protein mutations with support vector machines and evolutionary information. *Bioinformatics*. 2006;22(22):2729-34.
- [29] Thomas PD, Ebert D, Muruganujan A, Mushayahama T, Albu LP, Mi H. PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci*. 2022;31(1):08-22.
- [30] Choi Y, Chan AP. PROVEAN web server: A tool to predict the functional effect of amino acid substitutions and indels. *Bioinformatics*. 2015;31(16):2745-47.
- [31] Capriotti E, Calabrese R, Fariselli P, Martelli PL, Altman RB, Casadio R. WS-SNPs&GO: A web server for predicting the deleterious effect of human protein variants using functional annotation. *BMC Genomics*. 2013;14:01-07.
- [32] Pejaver V, Urresti J, Lugo-Martinez J, Pagel KA, Lin GN, Nam HJ, et al. Inferring the molecular and phenotypic impact of amino acid variants with MutPred2. *Nat Commun*. 2020;11(1):5918.
- [33] Capriotti E, Fariselli P, Casadio R. I-Mutant2.0: Predicting stability changes upon mutation from the protein sequence or structure. *Nucleic Acids Res*. 2005;33(suppl_2):W306-10.
- [34] Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, et al. ConSurf 2016: An improved methodology to estimate and visualize evolutionary conservation in macromolecules. *Nucleic Acids Res*. 2016;44(W1):W344-50.
- [35] Venselaar H, Te Beek TA, Kuipers RK, Hekkelman ML, Vriend G. Protein structure analysis of mutations causing inheritable diseases. An e-Science approach with life scientist friendly interfaces. *BMC Bioinformatics*. 2010;11:548.
- [36] Sherry ST, Ward M, Sirotkin K. dbSNP—database for single nucleotide polymorphisms and other classes of minor genetic variation. *Genome Res*. 1999;9(8):677-79.
- [37] UniProt Consortium. UniProt: The Universal protein knowledge base in 2023. *Nucleic Acids Res*. 2023;51(D1):D523-31.
- [38] Irfan M, Iqbal T, Hashmi S, Ghani U, Bhatti A. In silico prediction and functional analysis of nonsynonymous SNPs in human CTLA4 gene. *Sci Rep*. 2022;12(1):20441.
- [39] Falahi S, Karaji AG, Koohyanizadeh F, Rezaeiemanesh A, Salari F. A comprehensive in Silico analysis of the functional and structural impact of single nucleotide polymorphisms (SNPs) in the human IL-33 gene. *Comput Biol Chem*. 2021;94:107560.
- [40] Wang L, Yao Y, Si D. MMP-3 gene regulates the carcinogenesis and metabolic process of ovarian cancer, evidence from a Chinese population: Observational study and meta-analysis. *Medicine*. 2023;102(50):e36471.
- [41] Durmanova V, Javor J, Parnicka Z, Minarik G, Ocenasova A, Vaseckova B, et al. Impact of MMP2 rs243865 and MMP3 rs3025058 polymorphisms on clinical findings in Alzheimer's disease patients. *Mediators Inflamm*. 2021;2021(1):5573642.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Biochemistry, Government Medical College, Manjeri, Kerala, India.
2. Assistant Professor, Department of Biochemistry, Government Medical College, Thrissur, Kerala, India.
3. Professor, Department of Biochemistry, Government Medical College, Manjeri, Kerala, India.
4. Associate Professor, Department of Biochemistry, Government Medical College, Manjeri, Kerala, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Shajee S Nair,
Government Medical College Manjeri Malappuram, Manjeri-676121, Kerala, India.
E-mail: dmo2007@gmail.com

PLAGIARISM CHECKING METHODS: [\[Jain H et al.\]](#)

- Plagiarism X-checker: Jul 02, 2024
- Manual Googling: Aug 06, 2024
- iThenticate Software: Sep 24, 2024 (14%)

ETYMOLOGY: Author Origin

EMENDATIONS: 8

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? No
- For any images presented appropriate consent has been obtained from the subjects. No

Date of Submission: **Jul 01, 2024**
Date of Peer Review: **Aug 01, 2024**
Date of Acceptance: **Sep 25, 2024**
Date of Publishing: **Nov 01, 2024**