

Pattern of Refractive Error among Patients attending a Tertiary Care Hospital in Southern Assam, India: A Cross-sectional Study

NILANJAN KAUSHIK THAKUR¹, RUMA DAS DEBSIKDAR², SRAWANI SARKAR³, SURABHI PRASAD⁴, S LAKSHMI⁵, DEVAJIT DEKA⁶



ABSTRACT

Introduction: Uncorrected Refractive Errors (URE) are a primary cause of vision impairment worldwide and a leading contributor to blindness. These errors impact daily productivity, limit job opportunities and can significantly diminish quality of life, increasing both morbidity and financial burdens. Consequently, understanding the prevalence and types of UREs is essential for primary care physicians, ophthalmologists and policymakers.

Aim: To assess the pattern and magnitude of UREs among patients at a tertiary care hospital in South Assam, India.

Materials and Methods: A cross-sectional study was conducted at Silchar Medical College and Hospital, Silchar, Assam, India from May 2023 to April 2024, which included patients aged five years and older visiting the ophthalmology department. Patients with visual acuity worse than 6/6, which improved with pinhole testing, were further evaluated. Each participant underwent a comprehensive ophthalmic examination, including refraction testing. Data on uncorrected and corrected visual acuity, age, and sex were gathered. Spherical equivalents were used for refractive error analysis. The difference between categorical variables was estimated using the Chi-square test.

Results: Of the 11,932 participants screened, 6,112 were males (51.22%) and 5,820 were females (48.78%), aged 5-89 years. UREs were identified in 1,434 individuals (12%), with a median age of 41 years. In the URE group, 718 (50.1%) were females and 716 (49.9%) were males. The mean age of males was 39.51 ± 18.16 years, and that of females was 38.44 ± 19.12 years. The mean refractive error was -0.58 ± 3.11 DS. Emmetropia was present in 469 (32.7%), myopia in 427 (29.8%), high myopia in 38 (2.7%), and hypermetropia in 500 (34.9%). A total of 264 (18.4%) patients had astigmatism greater than 0.5. Anisometropia greater than 1.0 was present in 67 (4.7%). The proportion of different refractive errors was similar in males and females.

Conclusion: This study offers key insights into the magnitude and types of UREs in South Assam, India. The proportion of UREs was found to be 12%, with hypermetropia being the most common refractive error, followed by myopia. These findings underscore the need for targeted intervention programmes to improve access to corrective lenses and address vision impairment among these communities.

Keywords: Anisometropia, Astigmatism, Hypermetropia, Myopia, Uncorrected refractive error

INTRODUCTION

The UREs are the leading cause of moderate to severe vision impairment globally and the second most common cause of blindness. According to the findings in the Global Burden of Disease 2010 study [1], 101.2 million cases of Moderate and Severe Visual Impairment (MSVI) and 6.8 million cases of blindness were due to URE. A systematic review by Sheeladevi S et al., estimates a 53% prevalence of at least 0.50 D of SE ametropia (myopia 27.7%, hyperopia 22.9%) in India [2]. URE drastically reduces productivity, educational and job opportunities, and impacts the overall quality of life, leading to social isolation, increased morbidity, and economic distress for individuals and their families. Corrective glasses provide a low-cost and efficient solution in low-resource countries like India. However, there is a lack of data on the magnitude of the problem and suboptimal activity to address the issue, even though URE is part of the blindness control programme [3].

Refractive error is an established and significant public health problem. It is important for primary care physicians, ophthalmologists and policymakers to understand the magnitude and types of UREs among their patients. Targeting refractive error can have a huge impact on people's lives as well as improve the productivity of the country. In a randomised case-control research trial on 781 tea

pickers in Assam, presbyopic correction increased the daily weight of tea picking by 21.7% in the intervention group [4].

There have been only a few studies [5,6] attempting to understand this in the North-east part of India, and no study has been performed, to the best of our knowledge, to assess the prevalence of URE in southern Assam, India. The findings from this study are expected to provide future direction to policymakers and ophthalmologists in developing programmes to address this leading cause of vision impairment and to understand the existing barriers for accessing prescription glasses for correction. This study aimed to examine URE in patients and to determine the prevalence of URE in the patient population, the magnitude and pattern of refractive errors, and their variance across age groups and gender.

MATERIALS AND METHODS

A cross-sectional study was conducted at Silchar Medical College and Hospital in South Assam, India, between May 2023 and April 2024. The study adhered to the guidelines of the Declaration of Helsinki and was approved by the Silchar Medical College and Hospital Institutional Ethical Committee (IEC# SMC/5675). Informed consent was obtained from each participant before including them in the study.

Inclusion criteria: Patients attending the ophthalmology department, visual acuity worse than 6/6 that improved with a pinhole and patients with age >5 years were included in the study.

Exclusion criteria: Patients with any pathology in the lid, conjunctiva, cornea, anterior chamber, iris, lens, or posterior segment that impaired vision were excluded from the study.

Study Procedure

All patients attending the ophthalmology department at Silchar Medical College and Hospital (SMCH), which primarily serves underprivileged populations in southern Assam, India, were screened for visual acuity, and those aged >5 years with visual acuity worse than 6/6 that improved with a pinhole were selected. Distant visual acuity assessment was conducted using a standard illuminated Snellen visual acuity chart or a C-chart, both with and without a pinhole. Automated refraction followed by subjective refraction was performed on all patients for both distant and near vision. Objective refraction using a streak retinoscope (Beta 200; Heine Optotechnik, Germany) was also performed. Cycloplegic refraction using two drops of cyclopentolate 1% was done for children up to 18 years of age. A detailed examination of the anterior and posterior segments of all patients was performed. A proper proforma containing the information from each patient was maintained.

The refractive error of both eyes was converted to SE, which is the spherical power plus half the cylindrical power. Emmetropia was defined as SE between -0.50 and +0.50 Diopter Sphere (DS), myopia was considered when SE <-0.50 DS, and hyperopia was considered when SE >+0.50 DS. High myopia was defined as SE <-5.0 DS. Astigmatism was measured in negative cylindrical power, with the axis of the cylinder 15 degrees on either side of the horizontal meridian for With The Rule (WTR) astigmatism, and 15 degrees of the vertical meridian for Against The Rule (ATR) astigmatism. Oblique astigmatism was defined as occurring between 15-75° and 105-165° [6,7]. Anisometropia was defined as a difference of >1.0D in SE between the two eyes [7,8].

Visual acuity was measured in both eyes; however, to maintain consistency in methodology and outcome parameters for comparison with previous studies [6-8], statistical analysis was conducted on the right eye only, since the correlation between the SE of both eyes was very high (Pearson correlation 0.883).

STATISTICAL ANALYSIS

Data were collected in an Excel file, and all analyses were conducted using SAS 9.4 (SAS Institute, Inc., Cary, NC, USA). The difference between categorical variables was estimated using the Chi-square test, with p<0.05 considered statistically significant.


RESULTS

Out of the total 11,932 participants attending the Department of Ophthalmology who were screened in the study, refractive errors were identified in 1,434 (12.0%) patients. The age distribution ranged from 5 to 70 years, with a median age of 41 years. The distribution of patients according to various age groups and gender is provided in [Table/Fig-1]. Among the total study subjects, 716 (49.9%) were males, while the remaining 718 (50.1%) were females. The mean age of males was 39.51±18.16 years, and the mean age of females was 38.44±19.12 years. The mean refractive error was -0.58±3.11 DS.

A graphical representation of the distribution of SEs of the right eye of subjects is represented in [Table/Fig-2]. A total of 264 (18.4%) patients had astigmatism >0.5. Of these, 20% had WTR astigmatism, 26% had ATR astigmatism, and the rest had oblique astigmatism. Anisometropia >1.0 was present in 67 (4.7%) patients, more frequently in the younger age group of 20-39 years (49.25% of total anisometropia). The distribution of different types of refractive errors (in SE) is provided in [Table/Fig-3].

Age (in years)	Male, n (%)	Female, n (%)	Total, n (%)
5-19	53 (3.7)	63 (4.4)	116 (8.1)
20-29	90 (6.3)	98 (6.8)	188 (13.1)
30-39	230 (16.0)	235 (16.4)	465 (32.4)
40-49	215 (15.0)	220 (15.3)	435 (30.3)
50-59	70 (4.9)	53 (3.7)	123 (8.6)
>60	58 (4.0)	49 (3.4)	107 (7.5)
Total n (%)	716 (49.9)	718 (50.1)	1434 (100)

[Table/Fig-1]: Age and gender distribution.




[Table/Fig-2]: Frequency distribution of Spherical Equivalents (SE) of right eye.

Age (years)	Emmetropia n (%)	Myopia n (%)	High myopia n (%)	Hypermetropia n (%)	Astigmatism n (%)	Anisometropia n (%)
5-19	47 (3.3)	79 (5.5)	15 (1.1)	22 (1.5)	31 (2.2)	12 (0.8)
20-29	82 (5.7)	89 (6.2)	10 (0.7)	32 (2.2)	56 (3.9)	18 (1.3)
30-39	99 (6.9)	125 (8.7)	5 (0.4)	106 (7.4)	76 (5.3)	15 (1.0)
40-49	101 (7.0)	88 (6.1)	4 (0.3)	138 (9.6)	49 (3.4)	11 (0.8)
50-59	99 (6.9)	31 (2.2)	2 (0.1)	111 (7.7)	34 (2.4)	8 (0.6)
>60	41 (2.9)	15 (1.1)	2 (0.1)	91 (6.4)	18 (1.3)	3 (0.2)
Total n (%)	469 (32.7)	427 (29.8)	38 (2.7)	500 (34.9)	264 (18.4)	67 (4.7)

[Table/Fig-3]: Prevalence of refractive errors in various age groups.

The trend of different types of refractive errors with age is shown in [Table/Fig-4]. The proportion of myopia increased with age, with the highest frequency of presentation between 30-39 years, and then declined thereafter. Hypermetropia had the highest frequency of presentation at 40-49 years of age and decreased with further increases in age.



[Table/Fig-4]: Progression with the increase in age of different types of refractive errors.

In terms of gender distribution, the proportion of refractive errors was similar in males and females, as shown in [Table/Fig-5]. However,

high myopia was more common in males compared to females (63.2% vs 36.8%), although this difference was not statistically significant (p-value=0.09).

Refractive error type	Male n (%)	Female n (%)	Total n (%)	p-value*
Emmetropia	240 (51.2)	229 (48.8)	469 (100)	0.51
Myopia	210 (49.2)	217 (50.8)	427 (100)	0.71
High myopia	24 (63.2)	14 (36.8)	38 (100)	0.09
Hypermetropia	242 (48.4)	258 (51.6)	500 (100)	0.39
Astigmatism	133 (50.4)	131 (49.6)	264 (100)	0.87
Anisometropia	37 (55.2)	30 (44.8)	67 (100)	0.37

[Table/Fig-5]: Gender distribution of refractive errors.
*Chi-square test was used for the difference in proportion

DISCUSSION

Present study sheds light on the prevalence of refractive errors in the outpatient population of Northeast India, with a prevalence rate of 12%. Emmetropia was present in 32.7% of the participants. Among the refractive errors identified, hypermetropia was the most common (34.9%), followed by myopia (29.8%).

Astigmatism was present in 18.4% of the population, and anisometropia was observed in 4.7%. The pattern of refractive errors varied with age, with myopia being more common among younger individuals (30-39 years), while hypermetropia predominantly affected older adults (40-49 years). Additionally, the proportion of high myopia was 2.7%, an important finding given the associated risks of myopic macular degeneration and retinal detachment. The proportion of refractive errors was similar in males and females.

A summary comparing present study with other similar refractive error studies from various regions of India is provided in [Table/Fig-6] [2,5,6,9-11]. Hypermetropia was the most common refractive error in present study, particularly among the older age group. This aligns with the findings of Hashemi H et al., who noted that hypermetropia was more common in older adults due to the ageing process and loss of accommodative ability [12]. They found a prevalence of hypermetropia of 30.9% in adults, which was close to present finding of 34.9%. The Andhra Pradesh Eye Disease Study [9] reported a prevalence rate of 9.83% for hyperopia in urban Indian populations [Table/Fig-6], which was lower than what was observed in present study [2,5,6,9-11]. Present study prevalence rates were also higher than those reported by Nangia V et al., in Central India [10], who found a prevalence of 18% in their study of the rural population of Central India.

The prevalence of myopia in present study (29.8%) was consistent with trends observed in other Asian populations. Wu Q et al., reported a similar prevalence of 29.8% in China, highlighting the growing concern over myopia, especially among younger individuals [13]. Hashemi H et al., found a myopia prevalence of 4.9% in children and 26.5% in adults [12]. Sheeladevi S et al., in a systematic review on refractive errors in India, reported a similar finding of 27.7% prevalence of myopia [2]. Raju P et al., in their study of a rural South Indian population, observed the prevalence of myopia and high myopia to be 26.99% and 3.71%, respectively, which was similar to present study findings [11]. Bhutia KL et al., found a myopia prevalence of 31.1% in their study [5]. Present study finding of high myopia prevalence (2.7%) aligns with Holden BA et al., systematic review, which reported a similar prevalence of 2.7% [14]. They also predicted an increase in the number of cases of high myopia to 9.8% of the world population by 2050, highlighting the importance of addressing the growing population of individuals with high myopia. Studies have also shown an increasing trend in the prevalence of myopia and high myopia in the Asian population [15-17].

The prevalence of astigmatism in present study (18.4%) was slightly higher than the findings of Dandona R et al., in Southern

Author's name and year (Reference No.)	Place of study	Prevalence of refractive errors (%)	Most common refractive error type	Age group most affected
Dandona R et al., (1999) [9]	Urban population of South India	Not specified	≤15 years of age: Hyperopia 59.37% >15 years of age: Myopia 19.39%	≤15 years of age
Nangia V et al., (2010) [10]	Rural region in Central India	Myopia 31.3% Hyperopia 18.0%	Myopia 31.3% Hyperopia more common in lower age group	Not specified (Study population: >30 years)
Sheeladevi S et al., (2019) [2]	India (Systematic Review)	Prevalence of refractive error -53.1%	Myopia 27.7%	Not specified (Study population: >30 years)
Raju P et al., (2004) [11]	Rural South India	Myopia: 26.99%, high myopia: 3.71% Hyperopia: 18.70%	Myopia 26.99%	Not specified (Study population: >39 years)
Bhutia KL et al., (2021) [5]	Sikkim, India	6.7%	Myopia 31.1%	School going children
Natung T et al., (2017) [6]	Northeast India	55.56%	Myopia 27.4%	20-59 years (study population: ≥5 years)
Thakur NK et al., (2025)	South Assam, India	12%	Hypermetropia 34.9%	30-49 years (study population: ≥5 years)

[Table/Fig-6]: Comparison of present study with other refractive error studies conducted in different parts of India [2,5,6,9-11].

India [9], where 13% of the population had astigmatism, but lower than a population-based cohort study conducted in Bangladesh by Bourne RR et al., which reported a prevalence of 32.4% [7]. In a systematic review by Zhang J et al., the prevalence of astigmatism ranged between 8 to 62% [18]. They also noted that astigmatic patients experienced poor vision quality, increased glare and halos, difficulty with night driving, and a higher risk of falls. Vision-related tasks were performed more slowly by astigmatic patients compared with individuals who had appropriate glasses correction. The higher prevalence of astigmatism in our population highlights the need for early detection to prevent long-term visual impairment and poor productivity in vision-related tasks, especially in children.

Present study also found a lower proportion of anisometropia (4.7%) compared to studies by Bourne RR et al., which reported an anisometropia prevalence of around 8.8%. Another study by Zhou Y et al., found the prevalence of anisometropia to be 25.6%, which was higher than present study findings [7,19]. The lower rates in present study may be due to the hospital-based nature of the sample, which may not capture the full extent of anisometropia in the community. In a study by Hashemi H et al., the most common cause of amblyopia was anisometropia [20]. Hashemi H et al., emphasised the importance of early detection of anisometropia, particularly in younger populations, to prevent amblyopia and other vision-related problems. Yekta A et al., found that anisometropia-related amblyopia was present in 58.1% of their study subjects [21]. The prevalence of anisometropia in their study was 2.31%, which was lower than present study findings. The results of present study support the call for routine eye screenings, especially for children and adolescents, to detect anisometropia early and prevent its progression.

Refractive errors are the most prevalent cause of treatable visual impairment. They can be easily ascertained in everyday clinical practice quickly, easily, and safely. While refractive errors cannot be prevented, studies are still underway to understand how to do so; they can be easily treated with corrective glasses. If corrected in time, they can prevent the development of amblyopia and lifelong

visual impairment. It is pertinent that policies are developed to provide good quality and affordable refractive error corrective services and glasses to people belonging to low socio-economic status, especially in areas with limited eye care services. Our hospital caters to a diverse multicultural population, where a significant percentage lives in abject poverty (30-36%) according to a report by the Government of India [22]. As per our knowledge, no data on the prevalence or pattern of refractive errors exist from this area.

Limitation(s)

The study sample for this research was drawn from a hospital's OPD and not directly from the general population. Since the aim of the study was to focus only on URE, only patients with dimness of vision due to refractive error were included, while those with any pathology affecting the lid, conjunctiva, cornea, anterior chamber, iris, lens, or posterior segment that impaired vision were excluded from the study. However, many patients with temporary eye conditions, such as infective or allergic conjunctivitis, anterior uveitis, ocular foreign bodies, and corneal abrasions, who regularly visit our OPD, were also excluded. Such patients may have refractive errors but were excluded at the time of the study due to the presence of pathology at presentation.

Through this study, authors hope to provide insight into the magnitude and pattern of refractive error in the patient population and the need for correcting refractive errors at an early stage. Authors aim to provide evidence-based data to policymakers and non-governmental organisations so that resources and assistance can be allocated for correcting refractive errors, such as providing prescription glasses free of cost to patients. A substantial number of patients come from very low socio-economic backgrounds and struggle to meet their daily livelihoods, making them unwilling to spend money on corrective glasses.

CONCLUSION(S)

This study highlights the significant prevalence and distribution of refractive errors, particularly hypermetropia and myopia, in the outpatient population of South Assam in Northeast India. The findings emphasise the growing burden of myopia among younger individuals and the predominance of hypermetropia in older adults, which is consistent with global and regional trends. The notable prevalence of astigmatism and anisometropia further underscores the need for early detection and appropriate interventions to prevent long-term visual impairment. Present study results support the implementation of regular eye screenings, especially in underserved populations, to reduce the burden of UREs. Addressing these visual impairments through affordable corrective lenses and accessible eye care services is crucial for improving the quality of life in this region. Further population-based studies are necessary to provide a broader understanding of refractive error patterns and their public health impact.

REFERENCES

- [1] Lou L, Yao C, Jin Y, Perez V, Ye J. Global patterns in health burden of uncorrected refractive error. *Invest Ophthalmol Vis Sci*. 2016;57(14):6271-77.
- [2] Sheeladevi S, Seelam B, Nukella PB, Borah RR, Ali R, Keay L. Prevalence of refractive errors, uncorrected refractive error, and presbyopia in adults in India: A systematic review. *Indian J Ophthalmol*. 2019;67(5):583-92.
- [3] Holden B, Davis S, Jong M, Resnikoff S, editors. The evolution of uncorrected refractive error as a major public health issue. *Journal and Proceedings of the Royal Society of New South Wales*. 2014;147(453-454):101-06.
- [4] Reddy PA, Congdon N, MacKenzie G, Gogate P, Wen Q, Jan C, et al. Effect of providing near glasses on productivity among rural Indian tea workers with presbyopia (PROSPER): A randomised trial. *Lancet Glob Health*. 2018;6(9):e1019-e27.
- [5] Bhutia KL, Bhutia SC, Gupta N, Shenga DO. Prevalence of refractive errors among the school-going children in East Sikkim. *Indian J Ophthalmol*. 2021;69(8):2018-20.
- [6] Natung T, Taye T, Lyngdoh LA, Dkhar B, Hajong R. Refractive errors among patients attending the ophthalmology department of a medical college in North-East India. *J Family Med Prim Care*. 2017;6(3):543-48.
- [7] Bourne RR, Dineen BP, Ali SM, Noorul Huq DM, Johnson GJ. Prevalence of refractive error in Bangladeshi adults: Results of the National Blindness and Low Vision Survey of Bangladesh. *Ophthalmology*. 2004;111(6):1150-60.
- [8] Wong TY, Foster PJ, Hee J, Ng TP, Tsielsch JM, Chew SJ, et al. Prevalence and risk factors for refractive errors in adult Chinese in Singapore. *Invest Ophthalmol Vis Sci*. 2000;41(9):2486-94.
- [9] Dandona R, Dandona L, Naduvilath TJ, Srinivas M, McCarty CA, Rao GN. Refractive errors in an urban population in southern India: The Andhra Pradesh Eye Disease Study. *Invest Ophthalmol Vis Sci*. 1999;40(12):2810-18.
- [10] Nangia V, Jonas JB, Sinha A, Matin A, Kulkarni M. Refractive error in central India: The Central India Eye and Medical Study. *Ophthalmology*. 2010;117(4):693-99.
- [11] Raju P, Ve Ramesh S, Arvind H, George R, Baskaran M, Paul PG, et al. Prevalence of refractive errors in a rural south Indian population. *Invest Ophthalmol Vis Sci*. 2004;45(12):4268-72.
- [12] Hashemi H, Fotouhi A, Yekta A, Pakzad R, Ostadimoghaddam H, Khabazkhoob M. Global and regional estimates of prevalence of refractive errors: Systematic review and meta-analysis. *J Curr Ophthalmol*. 2018;30(1):03-22.
- [13] Wu Q, Tian Q, Zhang X, Xu J, Tang G, Li R, et al. Prevalence of refractive error and visual acuity among school children in the plateau region of Qinghai, China. *Int J Gen Med*. 2021;14:5795-805.
- [14] Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology*. 2016;123(5):1036-42.
- [15] Morgan IG, French AN, Ashby RS, Guo X, Ding X, He M, et al. The epidemics of myopia: Aetiology and prevention. *Prog Retin Eye Res*. 2018;62:134-49.
- [16] Morgan IG, Ohno-Matsui K, Saw SM. Myopia. *Lancet*. 2012;379(9827):1739-48.
- [17] GBD 2019 Blindness and Vision Impairment Collaborators; Vision Loss Expert Group of the Global Burden of Disease Study. Causes of blindness and vision impairment in 2020 and trends over 30 years, and prevalence of avoidable blindness in relation to VISION 2020: The Right to Sight: An analysis for the Global Burden of Disease Study. *Lancet Glob Health*. 2021;9(2):e144-e60.
- [18] Zhang J, Wu Y, Sharma B, Gupta R, Jawla S, Bullimore MA. Epidemiology and burden of astigmatism: A systematic literature review. *Optom Vis Sci*. 2023;100(3):218-31.
- [19] Zhou Y, Zhang XF, Chen XJ, Wang M, Cai JR, Xiong YJ, et al. Prevalence of anisometropia and influencing factors among school-age children in Nantong, China: A cross-sectional study. *Front Public Health*. 2023;11:1190285.
- [20] Hashemi H, Pakzad R, Yekta A, Bostamzad P, Aghamirsalim M, Sardari S, et al. Global and regional estimates of prevalence of amblyopia: A systematic review and meta-analysis. *Strabismus*. 2018;26(4):168-83.
- [21] Yekta A, Fotouhi A, Hashemi H, Dehghani C, Ostadimoghaddam H, Heravian J, et al. The prevalence of anisometropia, amblyopia and strabismus in schoolchildren of Shiraz, Iran. *Strabismus*. 2010;18(3):104-10.
- [22] NITI Aayog Gol. National Multidimensional Poverty Index, A progress review 2023. Report. [Internet]. [cited 2024 Apr 01]. Available from: <https://www.niti.gov.in/sites/default/files/2023-08/India-National-Multidimensional-Poverty-Index-2023.pdf>; 2023.

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of Ophthalmology, Silchar Medical College and Hospital, Silchar, Assam, India.
2. Professor, Department of Ophthalmology, Silchar Medical College and Hospital, Silchar, Assam, India.
3. Resident, Department of Ophthalmology, Silchar Medical College and Hospital, Silchar, Assam, India.
4. Resident, Department of Ophthalmology, Silchar Medical College and Hospital, Silchar, Assam, India.
5. Resident, Department of Ophthalmology, Silchar Medical College and Hospital, Silchar, Assam, India.
6. Resident, Department of Ophthalmology, Silchar Medical College and Hospital, Silchar, Assam, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Srawani Sarkar,
S-9, Resident Doctors Hostel, Silchar Medical College and Hospital,
Silchar-788014, Assam, India.
E-mail: srawani.sarkar@gmail.com

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Aug 05, 2024
- Manual Googling: Oct 24, 2024
- iThenticate Software: Nov 11, 2024 (13%)

ETYMOLOGY: Author Origin

EMENDATIONS: 8

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. NA

Date of Submission: **Aug 03, 2024**

Date of Peer Review: **Sep 16, 2024**

Date of Acceptance: **Nov 13, 2024**

Date of Publishing: **Jan 01, 2025**