

Insights into De Quervain's Tenosynovitis as an Unintended Consequence of Problematic Smartphone usage: A Cross-sectional Observational Study

ASWIN SUGUMARAN¹, DILIP KUMAR NAIDU², MADHAN RAJU³, SABARI VAASAN LAKSHMIKANTHAN⁴, S KARTHIKEYAN⁵, KEVIN DHAS⁶, VAMSI KRISHNA⁷, PRAVEEN RAJA⁸



ABSTRACT

Introduction: Smartphone use has been linked to Musculoskeletal Disorders (MSDs), particularly wrist and hand conditions like De Quervain's Tenosynovitis (DQT), which causes inflammation of the tendons of the thumb, mainly the Abductor Pollicis Longus (APL) and Extensor Pollicis Brevis (EPB). In India, the rise in smartphone usage has led to a corresponding increase in such disorders.

Aim: To investigate the association between problematic smartphone usage and the development of DQT and to assess its physical impact using the Visual Analogue Scale (VAS), Patient-rated Wrist/Hand Evaluation (PRWE) and quick Disabilities of the Arm, Shoulder and Hand (DASH) scores.

Materials and Methods: A cross-sectional observational study was conducted at the Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India, from July 2024 to September 2024, with 300 participants aged 20-50 years. Participants were classified based on the results of Finkelstein's Test and their smartphone usage patterns. Pain intensity was assessed using the VAS, while functional disability was measured using the quick DASH and PRWE questionnaires.

Results: The DQT was more prevalent in younger individuals (20-30 years) and females, with prolonged smartphone use (>8 hours/day) and frequent horizontal screen orientation strongly linked to its occurrence. Specific hand gestures such as Horizontal screen Both Thumb-Both Hand Grip (HBT-BHG) were common in bilateral DQT, while Horizontal screen Unilateral thumb-Isolated Hand Grip (HUT-IHG) was more frequent in unilateral cases. Vertical screen Unilateral Index finger-Other Hand Grip (VUI-OHG) and Vertical screen Unilateral Thumb-Other Hand Grip (VUT-OHG) gestures showed a lower incidence of DQT. Group 1 experienced higher pain intensity (VAS score 5.43 ± 1.34) and moderate disability scores (Quick DASH score 38.45 ± 12.52 ; PRWE score 42.67 ± 14.24), while Group 2 had minimal pain (VAS score 2.3 ± 1.02) and minimal disability scores (Quick DASH score 11.96 ± 6.68 and PRWE score 15.54 ± 8.46).

Conclusion: Problematic smartphone usage patterns (usage of phones for over 8 hours per day and horizontal screen orientation) were associated with the development of DQT, resulting in moderate disability scores that affected daily activities.

Keywords: Musculoskeletal diseases, Orthopaedics, Overuse injuries, Tendon entrapment

INTRODUCTION

The widespread use of smartphones has been a significant technological development of the 21st century. Excessive and incorrect use of smartphones may lead to Musculoskeletal Disorders (MSDs) that can impact the neck, shoulders, arms, hands and wrists [1]. One such condition is De Quervain's Tenosynovitis (DQT), also known as De Quervain's Disease (DQD). The increase in smartphone usage has been recognised as a major factor contributing to the rising incidence of DQT [2]. DQT is a painful condition affecting the tendons on the lateral side of the wrist, where patients may experience numbness, tingling, burning and cramping in addition to pain. The condition arises from inflammation of the tendons that control thumb movement, particularly the Abductor Pollicis Longus (APL) and Extensor Pollicis Brevis (EPB) [3]. These tendons pass through a narrow tunnel known as the first extensor compartment of the wrist, which is a fibro-osseous sheath that undergoes constriction and inflammation in cases of DQT [4].

The peak incidence of DQT occurs in individuals in their 40s and 50s and the prevalence is higher in women than in men [5]. Repeated thumb movements and prolonged awkward hand postures related to smartphone use, such as texting and gaming, have been linked to the onset of DQT [6]. Problematic smartphone use is associated

with the prevalence of DQT in empirical studies. For instance, a study of college students in Guangxi, China, found that smartphone use—particularly for games and social media—was linked to an increased risk of DQT [6]. Furthermore, research by Benites-Zapata VA et al., found a higher prevalence of DQT symptoms in individuals with problematic smartphone use [7].

Although the studies cited above have established an association between DQT and problematic phone usage, there has been no evaluation of smartphone handling techniques, such as horizontal versus vertical screen orientation and the various usage gestures. The present study aimed to elucidate the correlation between problematic smartphone usage and the gestures associated with using smartphones that are related to the development of DQT, as well as to assess its physical impact on daily activities using the VAS, Patient-rated Wrist/Hand Evaluation (PRWE) and Quick Disabilities of the Arm, Shoulder and Hand (DASH) scores.

MATERIALS AND METHODS

This was a cross-sectional observational study conducted at the Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India, from July 2024 to September 2024.


Approval from the Institutional Ethics Committee was exempted because it was a cross-sectional observational study.

Inclusion and Exclusion criteria: The present study included participants aged 20-50 years presenting to the Outpatient Department (OPD) with wrist and hand pain, who were willing to share their smartphone usage behaviour. Patients with a history of wrist injury, previous upper limb surgery, inflammatory joint disorders, congenital anomalies, pregnancy, lactation, a family history of DQT, or lateral/medial epicondylitis were excluded.


Sample size calculation: A sample size of 300 patients was classified into two groups based on the results of Finkelstein's Test. The sample size was calculated with an estimated prevalence of 50% for DQT, with 95% confidence and a 5% margin of error (Ali M et al., [8]).

Study Procedure

Patients meeting the inclusion criteria were included in the study. Participants were classified into two groups based on the results of Finkelstein's Test [9]: Group 1 (positive test, indicating DQT) and Group 2 (negative test, no DQT). A self-prepared questionnaire [ANNEXURE-1] was used to assess smartphone usage patterns. This questionnaire included details about daily smartphone usage hours, screen size, most frequently used activities (e.g., texting, gaming, browsing, scrolling) and the most commonly used screen orientation (horizontal or vertical) with gestures while using the smartphone [Table/Fig-1a-d,2a-d]. The collected data evaluated demographic



[Table/Fig-1]: Graphical image: Horizontal screen orientation commonly used gestures; a) Horizontal screen Both Thumb-Both Hand Grip (HBT-BHG); b) Horizontal screen Unilateral thumb-Isolated Hand Grip (HUT-IHG); c) Horizontal screen Unilateral Index finger-Other Hand Grip (HUI-OHG); d) Horizontal screen Unilateral Thumb-Other Hand Grip (HUT-OHG).



[Table/Fig-2]: Graphical image: Vertical screen orientation commonly used gestures: a) Vertical screen Both Thumb-Both Hand Grip (VBT-BHG); b) Vertical screen Unilateral thumb-Isolated Hand Grip (VUT-IHG); c) Vertical screen Unilateral Index finger-Other Hand Grip (VUI-OHG); d) Vertical screen Unilateral Thumb-Other Hand Grip (VUT-OHG).

information, usage patterns and gesture distribution. Additionally, patients were classified into unilateral and bilateral DQT based on clinical examination and Finkelstein's Test results. This comparison will help assess whether problematic smartphone usage patterns contribute differently to unilateral versus bilateral cases.

Pain intensity was assessed using the VAS [10]. For the purpose of the present study, scores were categorised as follows: 0 (no pain), 1-2 (mild pain), 3-4 (moderate pain), 5-7 (severe pain) and 8-10 (extreme pain). Functional disability of the upper limb was measured using the Quick DASH questionnaire score [11], which provides a score ranging from 0-100, with higher scores indicating greater disability. For the present study, the scores were divided into five categories: 0-20 (minimal disability); 21-40 (mild disability); 41-60 (moderate disability); 61-80 (severe disability); and 81-100 (extremely severe disability). Similarly, for better interpretation of results, the PRWE questionnaire score [12,13], which also provides a score of 0-100 with higher scores indicating greater disability, was divided into the following sub-scores: 0-24 (minimal pain and minimal disability); 25-49 (mild pain and mild disability); 50-74 (moderate pain and moderate disability); and 75-100 (severe pain and severe disability).

STATISTICAL ANALYSIS

The statistical analysis was performed using IBM Statistical Package for Social Sciences (SPSS) version 23.0. The qualitative variables were expressed as numbers and percentages. The Chi-square test was used to analyse the results, with a p-value of ≤ 0.05 considered statistically significant. The qualitative variables were described as mean \pm standard deviation. Analysis of variance was used to analyse qualitative variables.

RESULTS

A total of 300 participants were divided into two groups based on Finkelstein's Test, Group 1 (n=190) and Group 2 (n=110). Participants in Group 1 were predominantly aged 20-30 years (146 (76.8%)), whereas those in Group 2 were primarily aged 31-40 years (37 (33.64%)). A significant difference (p-value <0.0001*) was found between the age groups. Females (61.05%) had a greater incidence of DQT than males (p-value=0.0022*). Daily smartphone usage demonstrated a significant association with DQT, as 40% of DQT-positive individuals (n=76) used their phones for over eight hours per day, while the majority of DQT-negative individuals engaged in usage for less than four hours (p-value <0.0001*). Screen size and smartphone activities did not indicate any significant association with DQT. However, horizontal screen orientation was more prevalent among DQT-positive individuals (p-value=0.0168*) [Table/Fig-3].

Demographic characteristics and smartphone usage patterns		Group 1 (n=190)	Group 2 (n=110)	p-value
Age (in years)	20-30	146 (76.8%)	37 (33.64%)	<0.0001*
	31-40	38 (20%)	63 (57.27%)	
	41-50	6 (3.2%)	10 (9.09%)	
Gender (M/F)	Male	74 (38.95%)	63 (57.27%)	0.0022*
	Female	116 (61.05%)	47 (42.73%)	
Type of DQT	Unilateral	110 (57.89%)	-	-
	Bilateral	80 (42.11%)	-	
Daily usage (hours)	<4	49 (25.79%)	63 (57.27%)	<0.0001*
	4-8	65 (34.21%)	32 (29.09%)	
	>8	76 (40%)	15 (13.64%)	
Screen size (inches)	≤5.0	19 (10%)	17 (15.45%)	0.1810
	5.1-6.0	57 (30%)	38 (34.55%)	
	6.1-7.0	114 (60%)	55 (50%)	

Most used activity	Gaming	86 (45.26%)	41 (37.27%)	0.2095
	Texting	53 (27.89%)	32 (29.09%)	
	Scrolling	38 (20%)	22 (20%)	
	Videos	13 (6.84%)	15 (13.64%)	
Screen orientation	Horizontal	98 (51.58%)	41 (37.27%)	0.0168*
	Vertical	92 (48.42%)	69 (62.73%)	

[Table/Fig-3]: Demographic characteristics and smartphone usage patterns.

There were no significant differences between Group 1 and Group 2 in the use of HUT-OHG, HUI-OHG, HUT-IHG and HBT-BHG gestures. However, Group 2 showed that the VUT-OHG gesture was the second most commonly used gesture, indicating a lower incidence of DQT with this gesture. A highly significant difference was observed in the use of the VUI-OHG gesture, with Group 2 demonstrating it as the most commonly used gesture, which was associated with a lower incidence of DQT (p -value=0.0005). No significant differences were found between the two groups in the use of the VUT-IHG and VBT-BHG gestures [Table/Fig-4].

Gestures	Group 1 (n=190)	Group 2 (n=110)	p-value
HUT-OHG	11 (5.79%)	3 (2.73%)	0.2268
HUI-OHG	13 (6.84%)	6 (5.45%)	0.6343
HUT-IHG	29 (15.26%)	12 (10.91%)	0.2913
HBT-BHG	45 (23.68%)	20 (18.18%)	0.2659
VUT-OHG	11 (5.79%)	14 (12.73%)	0.0364*
VUI-OHG	17 (8.95%)	26 (23.64%)	0.0005*
VUT-IHG	29 (15.26%)	14 (12.73%)	0.5474
VBT-BHG	35 (18.42%)	17 (15.45%)	0.5132

[Table/Fig-4]: Comparison of common hand gestures usage between Group 1 (Finkelstein Test positive) and Group 2 (Finkelstein Test negative) (N=300).

HBT-BHG: Horizontal screen both thumb-both hand grip; HUT-IHG: Horizontal screen unilateral thumb-isolated hand grip; HUI-OHG: Horizontal screen unilateral index finger-other hand grip; HUT-OHG: Horizontal screen unilateral thumb-other hand grip; VBT-BHG: Vertical screen both thumb-both hand grip; VUT-IHG: Vertical screen unilateral thumb-isolated hand grip; VUI-OHG: Vertical screen unilateral index finger-other hand grip; VUT-OHG: Vertical screen unilateral thumb-other hand grip

Unilateral DQT was most commonly associated with the HUT-IHG gesture, followed by the VUT-IHG and HUI-OHG gestures, with these differences being statistically significant (p -value <0.05). For bilateral DQT, the most common gesture was HBT-BHG, followed by VBT-BHG, both showing statistically significant differences. No significant differences were found between unilateral and bilateral DQT in the use of the HUT-OHG, VUT-OHG and VUI-OHG gestures [Table/Fig-5].

Gestures Finkelstein's test positive (n=190)	Unilateral DQT (n=110)	Bilateral DQT (n=80)	p-value
HUT-OHG	8 (7.27%)	3 (3.75%)	0.3062
HUI-OHG	11 (10%)	2 (2.5%)	0.0438*
HUT-IHG	26 (23.64%)	3 (3.75%)	0.0002*
HBT-BHG	11 (10%)	34 (42.5%)	<0.0001*
VUT-OHG	6 (5.45%)	5 (6.25%)	0.8161
VUI-OHG	10 (9.09%)	7 (8.75%)	0.9356
VUT-IHG	24 (21.82%)	5 (6.25%)	0.0033*
VBT-BHG	14 (12.73%)	21 (26.25%)	0.0179*

[Table/Fig-5]: Comparison of hand gestures between unilateral and bilateral DQT.

HBT-BHG: Horizontal screen both thumb-both hand grip; HUT-IHG: Horizontal screen unilateral thumb-isolated hand grip; HUI-OHG: Horizontal screen unilateral index finger-other hand grip; HUT-OHG: Horizontal screen unilateral thumb-other hand grip; VBT-BHG: Vertical screen both thumb-both hand grip; VUT-IHG: Vertical screen unilateral thumb-isolated hand grip; VUI-OHG: Vertical screen unilateral index finger-other hand grip; VUT-OHG: Vertical screen unilateral thumb-other hand grip

In Group 1, the HUT-OHG gesture was used by 11 participants (5.79%), resulting in mild disability scores. The VUT-IHG gesture was used by 29 participants (15.26%) and the VBT-BHG gesture was used by 35 participants (18.42%), both of which were associated with severe pain and moderate disability scores [Table/Fig-6].

Gestures	Group 1 (n=190)	VAS	PRWE	Quick DASH
HUT-OHG	11 (5.79%)	4.36±0.67	46.64±6.76	36.55±7.63
HUI-OHG	13 (6.84%)	3.92±0.95	40.46±7.22	31.38±7.52
HUT-IHG	29 (15.26%)	5±0.85	50.83±5.64	40.93±9.54
HBT-BHG	45 (23.68%)	6.98±0.87	65.49±5.68	55.73±5.82
VUT-OHG	11 (5.79%)	4.09±0.83	29.64±5.35	24.63±3.5
VUI-OHG	17 (8.95%)	4.05±0.83	26.94±2.61	22.94±2.05
VUT-IHG	29 (15.26%)	5.22±0.83	55.17±10.7	45±5.66
VBT-BHG	35 (18.42%)	5.97±0.75	59.94±8.44	50.31±9.44

[Table/Fig-6]: Assessment of VAS, PRWE and Quick-DASH scores in participants with DQT (Group 1).

VAS: Visual analogue scale; PRWE: Patient-rated wrist/hand evaluation; Quick DASH: Disabilities of the arm, shoulder and hand; HBT-BHG: Horizontal screen both thumb-both hand grip;

HUT-IHG: Horizontal screen unilateral thumb-isolated hand grip; HUI-OHG: Horizontal screen unilateral index finger-other hand grip; HUT-OHG: Horizontal screen unilateral thumb-other hand grip;

VBT-BHG: Vertical screen both thumb-both hand grip; VUT-IHG: Vertical screen unilateral thumb-isolated hand grip; VUI-OHG: Vertical screen unilateral index finger-other hand grip;

VUT-OHG: Vertical screen unilateral thumb-other hand grip

In Group 2, the HUT-OHG gesture was used by 3 participants (2.73%), resulting in mild pain and minimal disability scores. Similarly, the HUI-OHG gesture was used by 6 participants (5.45%) and exhibited mild pain and minimal disability scores. The VUT-OHG gesture was used by 14 participants (12.73%) and the VUI-OHG gesture was used by 26 participants (23.64%), both displaying mild pain and minimal disability scores [Table/Fig-7].

Gestures	Group 2 (n=110)	VAS	PRWE	Quick-DASH
HUT-OHG	3 (2.73%)	1.02±0.32	14.43±4.24	11.67±3.56
HUI-OHG	6 (5.45%)	1.09±0.29	12.28±3.56	10.33±2.88
HUT-IHG	12 (10.91%)	2.21±1.41	16.06±6.82	12.55±3.12
HBT-BHG	20 (18.18%)	3.12±1.18	22.82±5.66	18.62±4.36
VUT-OHG	14 (12.73%)	1.13±0.03	10.55±4.86	9.88±1.98
VUI-OHG	26 (23.64%)	1.07±0.12	8.37±4.33	5.74±1.04
VUT-IHG	14 (12.73%)	3.14±1.13	18.54±5.34	14.48±2.84
VBT-BHG	17 (15.45%)	3.18±1.43	20.64±6.35	16.32±4.6

[Table/Fig-7]: Assessment of VAS, PRWE and Quick-DASH scores in participants without DQT (Group 2).

VAS: Visual analogue scale; PRWE: Patient-rated wrist/hand evaluation; Quick DASH: Disabilities of the arm, shoulder and hand; HBT-BHG: Horizontal screen both thumb-both hand grip;

HUT-IHG: Horizontal screen unilateral thumb-isolated hand grip; HUI-OHG: Horizontal screen unilateral index finger-other hand grip; HUT-OHG: Horizontal screen unilateral thumb-other hand grip;

VBT-BHG: Vertical screen both thumb-both hand grip; VUT-IHG: Vertical screen unilateral thumb-isolated hand grip; VUI-OHG: Vertical screen unilateral index finger-other hand grip;

VUT-OHG: Vertical screen unilateral thumb-other hand grip

Group 1 reported a significantly higher mean pain intensity VAS score of 5.43±1.34 (severe pain) compared to Group 2's score of 2.3±1.02 (mild pain) (p -value <0.0001). The mean PRWE score for Group 1 was 42.67±14.24 (mild disability), which was significantly higher than Group 2's score of 15.54±8.46 (minimal disability), indicating greater disability in Group 1, which is statistically significant (p -value <0.0001*). Similarly, Group 1 had a Quick DASH score of 38.45±12.52 (mild disability), compared to Group 2's score of 11.96±6.68 (minimal disability), which is statistically significant (p -value <0.0001*) [Table/Fig-8].

Physical impact score	Group 1 (n=190)	Group 2 (n=110)	p-value
VAS score	5.43±1.34	2.3±1.02	<0.0001*
PRWE score	42.67±14.24	15.54±8.46	<0.0001*
Quick-DASH score	38.45±12.52	11.96±6.68	<0.0001*

[Table/Fig-8]: Comparison of VAS, PRWE and Quick-DASH scores between Group 1 and Group 2.

VAS: Visual analogue scale; PRWE: Patient-rated wrist/hand evaluation; Quick DASH: Disabilities of the arm, shoulder and hand

DISCUSSION

In the present study, it was found that using phones for over eight hours per day and employing a horizontal screen orientation were

associated with the development of DQT, resulting in moderate disability scores that affected daily activities. The HBT-BHG gesture was the most commonly used in Group 1 (Finkelstein test-positive), while the VUI-OHG gesture was the most commonly used in Group 2 (Finkelstein test-negative). The HUT-IHG and VUT-IHG gestures were common in Unilateral DQT. The HBT-BHG and VBT-BHG gestures were the most commonly used in Bilateral DQT. Participants in Group 1 (Finkelstein test-positive) experienced higher pain intensity (VAS score) and moderate disability (Quick DASH and PRWE scores), whereas Group 2 reported minimal pain (VAS score) and minimal disability (Quick DASH and PRWE scores).

Berites-Zapata VA et al., found that problematic phone usage increases the risk of DQT in young adults [7]. This aligns with the observation that the majority of participants affected by DQT in the present study were aged between 20 and 30 years. Wolf JM et al., reported that women had a considerably elevated incidence of DQT, with an incidence of 2.8 per 1,000 persons/year, while men had an incidence of 0.6 per 1,000 persons/year [14]. Similarly, the present study revealed a higher proportion of females in the Finkelstein test-positive group (61.05%). They also noted that the incidence of DQT increases with age, particularly in individuals over 40 years old [14]. Morshed MK et al., suggested that extended daily smartphone usage, especially in younger age groups and the use of smaller-screen smartphones may increase the incidence of DQT in young people [15]. The present study revealed that excessive smartphone use (>8 hours daily), particularly involving repetitive thumb movements such as gaming and texting, significantly increases the risk of DQT. In contrast, screen size did not show a significant association with DQT. Ma T et al., also found that mobile gaming frequency, duration and wrist posture increased the DQT risk [16].

Similar to the present study, Blackburn J et al., found that DQT patients experienced greater pain and functional limitations [17]. Participants with DQT in the present study reported a significantly greater physical impact of pain, disability and functional limitations.

When assessing the impact of gestures, it was found that the HUT-IHG and HBT-BHG gestures were associated with higher pain, disability and functional limitations. In contrast, the VUT-OHG and VUI-OHG gestures were associated with lower pain and disability scores. The present study also revealed that prolonged use and frequent horizontal screen orientation may contribute to the development or severity of DQT.

Limitation(s)

The cross-sectional design of the present study makes it difficult to establish a causal relationship between smartphone use and DQT. Additionally, self-reported data may introduce bias or inconsistencies in the reporting of smartphone usage and pain levels.

CONCLUSION(S)

Based on the findings, it can be concluded that younger individuals (aged 20-30 years) and females had a higher prevalence of DQT. Prolonged smartphone usage (>8 hours/day) and frequent horizontal screen orientation were significantly associated with DQT. Specific hand gestures, such as HBT-BHG, were more commonly observed in individuals with DQT, while the VUI-OHG and VUT-OHG

gestures had a lower incidence of DQT. The HBT-BHG gesture was the most common in bilateral DQT, whereas the HUT-IHG gesture was the most frequently used in unilateral DQT. Subjects with DQT exhibited significantly higher pain intensity, with the mean PRWE score indicating moderate disability and the mean Quick DASH score also indicating moderate disability compared to participants without DQT. Thus, in this digital era, problematic phone usage has a strong association with the development of DQT. Further long-term follow-up multicentre studies with larger populations are needed to explore the multifactorial associations of DQT in relation to problematic smartphone usage.

REFERENCES

- [1] Mersal FA, Negm LM, Fawzy MS, Rajennal AT, Alanazi RS, Alanazi LO. Effect of mobile phone use on musculoskeletal complaints: Insights from nursing students at Northern Border University, Arar, Saudi Arabia. *Cureus*. 2024;16(3):e57181. Doi: 10.7759/cureus.57181.
- [2] Zirek E, Mustafaoglu R, Yasaci Z, Griffiths MD. A systematic review of musculoskeletal complaints, symptoms, and pathologies related to mobile phone usage. *Musculoskelet Sci Pract*. 2020;49:102196. Doi: 10.1016/j.msksp.2020.102196.
- [3] Fakoya AO, Tarzian M, Sabater EL, Burgos DM, Marty Gl. de Quervain's disease: A discourse on etiology, diagnosis, and treatment. *Cureus*. 2023;15(4):e38079. Doi: 10.7759/cureus.38079.
- [4] Chaya B, Bakhach E, Bakhach J. The de-Quervain tenosynovitis: Literature review. *Biomedical Journal*. 2018;8(4):6650-52. Doi: 10.26717/BJSTR.2018.08.001691.
- [5] Satteson E, Tannan SC. De Quervain Tenosynovitis. [Updated 2023 Nov 22]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK442005/>.
- [6] Nie X, Huang L, Hou J, Dai A, He L, Zheng P, et al. Smartphone usage behaviours and their association with De Quervain's Tenosynovitis (DQT) among college students: A cross-sectional study in Guangxi, China. *BMC Public Health*. 2023;23(1):2257. Doi: 10.1186/s12889-023-16808-z.
- [7] Benites-Zapata VA, Jiménez-Torres VE, Ayala-Roldán MP. Problematic smartphone use is associated with de Quervain's tenosynovitis symptomatology among young adults. *Musculoskelet Sci Pract*. 2021;53:102356. Doi: 10.1016/j.msksp.2021.102356.
- [8] Ali M, Asim M, Danish SH, Ahmad F, Iqbal A, Hasan SD. Frequency of De Quervain's tenosynovitis and its association with SMS texting. *Muscles Ligaments Tendons J*. 2014;4(1):74. Doi: 10.11138/mltj/2014.4.1.074.
- [9] Dawson C, Mudgal CS. Staged description of the Finkelstein test. *J Hand Surg Am*. 2010;35(9):1513-15. Doi: 10.1016/j.jhsa.2010.05.022.
- [10] Boomstra AM, Preuper HR, Reneman MF, Posthumus JB, Stewart RE. Reliability and validity of the visual analogue scale for disability in patients with chronic musculoskeletal pain. *Int J Rehabil Res*. 2008;31(2):165-69. Doi: 10.1097/MRR.0b013e3282fc0f93.
- [11] Gummesson C, Ward MM, Atroshi I. The shortened disabilities of the arm, shoulder and hand questionnaire (Quick DASH): Validity and reliability based on responses within the full-length DASH. *BMC Musculoskelet Disord*. 2006;7:01-07. Doi: 10.1186/1471-2474-7-44.
- [12] MacDermid JC, Turgeon T, Richards RS, Beadle M, Roth JH. Patient rating of wrist pain and disability: A reliable and valid measurement tool. *J Orthop Trauma*. 1998;12(8):577-86. Doi: 10.1097/00005131-199811000-00009.
- [13] MacDermid JC. The patient-rated wrist evaluation (PRWE)® user manual. Hamilton: McMaster University. 2007 Dec.
- [14] Wolf JM, Sturdivant RX, Owens BD. Incidence of de Quervain's tenosynovitis in a young, active population. *J Hand Surg Am*. 2009;34(1):112-15. Doi: 10.1016/j.jhsa.2008.08.020.
- [15] Morshed MK, Liton MS, Hossain MS. Relationship between overuse of smartphones and De Quervain's tenosynovitis in young adults. *The Planet*. 2023;7(01):289-95. Available from: <https://bdjournals.org/index.php/planet/article/view/431>.
- [16] Ma T, Song L, Ning S, Wang H, Zhang G, Wu Z. Relationship between the incidence of de Quervain's disease among teenagers and mobile gaming. *Int Orthop*. 2019;43:2587-92. Doi: 10.1007/s00264-019-04389-9.
- [17] Blackburn J, van der Oest MJ, Selles RW, Chen NC, Feitz R, Vranceanu AM, et al. Which psychological variables are associated with pain and function before surgery for de Quervain's tenosynovitis? A cross-sectional study. *Clin Orthop Relat Res*. 2019;477(12):2750-58. Doi: 10.1097/CORR.0000000000000992.

PARTICULARS OF CONTRIBUTORS:

- Junior Resident, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Professor, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Senior Resident, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Associate Professor, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Assistant Professor, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Assistant Professor, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Senior Resident, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.
- Senior Resident, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu, Tamil Nadu, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Aswin Sugumaran,
Junior Resident, Department of Orthopaedics, SRM Medical College Hospital and Research Centre, SRM Institute of Science and Technology, Chengalpattu-603203, Tamil Nadu, India.
E-mail: as1446@srmist.edu.in

PLAGIARISM CHECKING METHODS: [\[Jain H et al.\]](#)

- Plagiarism X-checker: Feb 01, 2025
- Manual Googling: Feb 25, 2025
- iThenticate Software: Mar 19, 2025 (9%)

ETYMOLOGY: Author Origin**EMENDATIONS:** 8**AUTHOR DECLARATION:**

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? No
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: **Jan 30, 2025**Date of Peer Review: **Feb 22, 2025**Date of Acceptance: **Mar 21, 2025**Date of Publishing: **Apr 01, 2025****[ANNEXURE-1]****Demographic, Smartphone usage patterns and Gesture Assessment Questionnaire:****Patient Information:**

- Age:
- Gender: Male Female

Wrist Health and Symptoms:

- Have you experienced wrist pain lasting for more than 3 weeks? Yes No
- Which wrist/hand is affected by the symptoms? Unilateral (One wrist/hand) Bilateral (Both wrists/hands)
- Finkelstein Test Result:** Positive Negative
- If the Finkelstein Test was Positive, please specify which side(s) were affected:**
 - Unilateral (One wrist/hand)
 - Bilateral (Both wrist/hands)

Smartphone Usage Patterns:

- How many hours per day do you typically use your smartphone?
 - Less than 4 hours
 - 4-8 hours
 - More than 8 hours
- What is the screen size of your smartphone?**
 - Less than 5 inches
 - 5.1-6.0 inches
 - 6.1-7.0 inches
- What activity do you most commonly do on your smartphone?**
 - Gaming
 - Texting

- Scrolling (e.g., social media)
- Watching videos

Smartphone Grip and Usage Gestures:

- Which screen orientation do you primarily use while interacting with your smartphone?
 - Horizontal
 - Vertical
- If you most commonly use a Horizontal screen orientation, what is your most commonly used gesture?
 - HBT-BHG: Horizontal screen, Both Thumbs – Both Hand Grip
 - HUT-IHG: Horizontal screen, Unilateral Thumb – Isolated Hand Grip
 - HUI-OHG: Horizontal screen, Unilateral Index Finger – Other Hand Grip
 - HUT-OHG: Horizontal screen, Unilateral Thumb – Other Hand Grip
- If you most commonly use a Vertical screen orientation, what is your most commonly used gesture?
 - VBT-BHG: Vertical screen, Both Thumbs – Both Hand Grip
 - VUT-IHG: Vertical screen, Unilateral Thumb – Isolated Hand Grip
 - VUI-OHG: Vertical screen, Unilateral Index Finger – Other Hand Grip
 - VUT-OHG: Vertical screen, Unilateral Thumb – Other Hand Grip