Original Article

Obstetrics and Gynaecology Section

Urine Dipstick versus Urine Protein Creatinine Ratio for the Diagnosis of Preeclampsia: A Cross-sectional Study

SUHAS GAIKWAD1, VIDYA GAIKWAD2, PRAGYA TIWARI3



ABSTRACT

Introduction: Preeclampsia is a condition related to pregnancy, manifesting after 20 weeks of gestation, characterised by hypertension and proteinuria, with potentially severe consequences for both the mother and the foetus.

Aim: To evaluate the diagnostic performance of the urine dipstick test and the Urine Protein-Creatinine Ratio (UPCR) against the 24-hour urine protein test as a reference for detecting proteinuria and subsequently, preeclampsia.

Materials and Methods: This was a cross-sectional study conducted at the Department of Obstetrics and Gynaecology at Dr. D. Y. Patil Medical College and Research Centre, Pune, Maharashtra, India, from October 2023 to October 2024. A total of 80 hypertensive pregnant women beyond 20 weeks of gestation were included. Participants were screened using the urine dipstick test and UPCR, followed by a 24-hour urine collection for protein estimation. Sensitivity, specificity, Positive Predictive

Value (PPV), Negative Predictive Value (NPV), accuracy and F1-score were calculated for both tests.

Results: The mean age of the study population was 27±5 years. The study included 35 (44%) cases of preeclampsia and 45 (56%) cases of gestational hypertension. Preeclampsia was more prevalent in patients aged 25-35 years, accounting for 17 (49%) cases. The urine dipstick test showed a sensitivity of 43%, specificity of 40%, PPV of 36%, NPV of 47%, accuracy of 41% and F1-score of 39%. The UPCR demonstrated a sensitivity of 91%, specificity of 7%, PPV of 43%, NPV of 50%, accuracy of 44% and F1-score of 59%.

Conclusion: The UPCR is more reliable for confirming proteinuria but has a low NPV, making it less effective at ruling out preeclampsia. These findings suggest that clinical assessment, combined with UPCR, should be used as an adjunct to improve early detection of proteinuria in preeclampsia.

Keywords: 24-hour urine protein test, Diagnostic performance, Hypertension, Pregnancy

INTRODUCTION

Preeclampsia is a condition related to pregnancy and is characterised by new-onset hypertension after 20 weeks of gestation [1]. The global incidence is estimated to be between 2% and 10%, while studies in India have reported a prevalence of 8% to 10% [2-4]. A typical diagnosis includes blood pressure of 140 mmHg or higher (systolic) or blood pressure of 90 mmHg or higher (diastolic) measured on two occasions at least four hours apart [1,5,6]. In addition to hypertension, proteinuria is a standard biomarker of preeclampsia [7-9]. The leakage of protein in urine is attributed to damage in the glomeruli as a result of hypertension [10]. Therefore, the presence of proteinuria alongside hypertension is a strong indication of preeclampsia. Identification and early management of preeclampsia are crucial for maternal and fetal outcomes.

The UPCR has been widely used as an alternative to the 24-hour urine collection, offering the advantage of being quicker, more convenient and readily available [11-13]. By calculating the ratio of protein to creatinine in a single urine sample, UPCR can efficiently identify proteinuria without the need for prolonged urine collection. Additionally, urine protein dipsticks are extensively used for initial screening, providing a rapid, though less precise, estimate of proteinuria [14,15]. Given the life-threatening nature of preeclampsia, which can result in end-organ failure and severe complications for both the mother and foetus, the development of reliable, accurate and time-efficient diagnostic tools like UPCR and urine dipsticks is essential for improving outcomes through early detection and timely intervention [16,17]. However, there is a dearth of knowledge regarding the reliability and specificity of

these time-saving diagnostic tools to ensure timely and appropriate management.

Here, the authors aimed to contribute to the existing literature by evaluating the diagnostic performance of the urine dipstick test and the UPCR against the 24-hour urine protein test.

MATERIALS AND METHODS

A cross-sectional study was conducted at the Department of Obstetrics and Gynaecology at Dr. D. Y. Patil Medical College and Research Centre, Pune, Maharashtra, India, from October 2023 to October 2024. The Institutional Ethics Committee approved the study (IESC/FP/2023/54) and written informed consent was obtained from all participants.

Sample size calculation: A total of 80 patients were included in the study. The sample size was calculated using Fisher's formula: $n=Z^2\times P$ (1-P)/e², where Z=1.96 for a 95% confidence interval, P=0.05 (prevalence of preeclampsia) [18,19] and e=0.05 (precision). With a 5% attrition rate, the sample was calculated to be 77.

Inclusion criteria: The study included hypertensive pregnant women with blood pressure levels of ≥140/90 mmHg, who had a singleton pregnancy and were beyond 20 weeks of gestation.

Exclusion criteria: Women with known renal diseases, urinary tract infections, dehydration, chronic hypertension, diabetes mellitus, molar pregnancy, or multiple pregnancies were excluded. Additionally, patients who delivered before the collection of the 24-hour urine sample were also excluded. These conditions were excluded due to their potential to alter proteinuria levels, which could confound the results and affect the outcomes of the study.

Study Procedure

Laboratory analysis: After obtaining informed consent, demographic details and obstetric parameters, including age and gravidity, were recorded. A random spot urine sample for dipstick and UPCR testing was taken from each participant, followed by the collection of 24-hour urine output for the estimation of total protein.

Urine dipstick test: The urine sample was tested using commercially available urine dipsticks (Siemens Uristix reagent strips). The dipstick results were interpreted as negative, trace, 1+, 2+, or 3+ and values of $\geq 1+$ were considered positive for proteinuria.

Urine Protein-Creatinine Ratio (UPCR): The urine sample was analysed for the estimation of protein and creatinine concentrations. The Urinary Protein-to-Creatinine Ratio (UPCR) was calculated as (urinary protein/urinary creatinine), with a threshold value of \geq 0.3 considered positive for significant proteinuria [20]. Protein was measured using the pyragallol method and creatinine levels were determined using the Jaffe method [21].

24-Hour Urine Protein Test (Gold Standard): Urine output over 24 hours was collected in a sterile container and analysed for the estimation of urine protein. Total protein excretion was measured using an automated biochemical analyser, with ≥300 mg/day considered indicative of proteinuria.

STATISTICAL ANALYSIS


Data were analysed using GraphPad Prism 10. Sensitivity, specificity, PPV, NPV, accuracy and F1-score were calculated for both the urine dipstick and UPCR, using the 24-hour urine protein test as the reference standard. Categorical variables were expressed as frequencies and percentages.

RESULTS

The study population consisted of 80 hypertensive patients, of whom 35 (44%) had preeclampsia and 45 (56%) had gestational hypertension. The age distribution of the study population showed that the majority of cases were in the 20-25 years age group 38 (48%). The least represented age group was 18-20 years 2 (3%) [Table/Fig-1]. The majority of the patients with preeclampsia were within the age group of 25-30 years [Table/Fig-2].

Age group (in years)	n (%)
18-20	2 (2.5)
20-25	38 (47.5)
25-30	21 (26.25)
30-35	15 (18.75)
35-40	4 (5)
Total	80 (100)

[Table/Fig-1]: Distribution of population based on age.



The highest proportion of participants had no prior pregnancies 25 (31%), followed by those with a gravidity of 2 24 (30%) [Table/Fig-3].

Gravidity	n (%)	
0	25 (31.25)	
1	2 (2.5)	
2	24 (30)	
3	14 (17.5)	
>3	15 (18.75)	
Total	80 (100)	
[Table/Fig-3]: Distribution of population based on gravidity.		

The diagnostic performance of the urine dipstick test was evaluated against the 24-hour urine protein test (the gold standard) in patients. Among the 80 patients, the urine dipstick test was positive in 42 cases, of which 15 (36%) were confirmed positive by the 24-hour urine protein test, while 27 (64%) were false positives [Table/Fig-4].

Preeclampsia				
Urine dipstick	Positive	Negative	Total	
Positive	15	27	42	
Negative	20	18	38	
Total	35	45	80	
[Table/Fig. 4]. Diagnostic performance of using direction text for procedure in				

Furthermore, the diagnostic performance of the urine UPCR was evaluated against the 24-hour urine protein test in patients. Among the 80 patients, UPCR was positive in 74 cases, of which 32 (43%) were confirmed positive by the 24-hour urine protein test, while 42 (57%) were false positives [Table/Fig-5].

Preeclampsia				
Urine Protein-Creatinine Ratio (UPCR)	Positive	Negative	Total	
Positive	32	42	74	
Negative	3	3	6	
Total	35	45	80	
[Table/Fig-5]: Diagnostic performance of UPCR for preeclampsia.				

The sensitivity of the dipstick was 43%, while UPCR had a sensitivity of 91%. The PPV for the dipstick was 36%, while UPCR showed a higher PPV of 43%, demonstrating its better ability to confirm positive cases. The NPV was low for the dipstick, with values of 47% for the dipstick and 50% for UPCR. The F1-score, which reflects the balance between precision and recall, was 39% for the dipstick and 59% for UPCR [Table/Fig-6].

Metric	Dipstick (95% CI)	UPCR (95% CI)
Sensitivity	43% (31.51%, 54.21%)	91% (73.67%, 109.19%)
Specificity	40% (28.71%, 51.29%)	7% (0.75%, 12.60%)
PPV	36% (24.42%, 46.99%)	43% (31.84%, 54.64%)
NPV	47% (36.08%, 58.65%)	50% (38.73%, 61.27%)
Accuracy	41% (29.96%, 52.54%)	44% (32.50%, 55.00%)
F1-Score	39% (17.51%, 60.47%)	59% (42.21%, 75.73%)

[Table/Fig-6]: Comparison of urine dipstick and Urine Protein-to-Creatinine Ratio (UPCR) for detecting preeclampsia.

DISCUSSION

The present study evaluated the diagnostic performance of the urine dipstick test and the UPCR against the 24-hour urine protein test. The age distribution analysis showed that the majority of cases (48%) were within the age range of 20-25 years. Further analysis of the population based on the presence and absence of preeclampsia revealed that, for those aged over 25 years, there were more incidences of preeclampsia. These findings align with previous epidemiological studies indicating that preeclampsia predominantly affects women in this age group. A systematic review by Duckitt K and Harrington D shows that pregnant women aged over 30 years

exhibit a gradual increase in the odds of developing preeclampsia [22]. Further studies by Chan TF et al., and Sheen JJ et al., are in agreement with our findings [23,24].

The urine dipstick test demonstrated a sensitivity of 43% and specificity of 40% in diagnosing preeclampsia, suggesting that it missed a significant number of cases. A meta-analysis study by Teeuw HM et al., analysing 19 studies (n=3700 urine samples), showed that the urine dipstick does not efficiently exclude preeclampsia patients from those with gestational hypertension [25]. Phelan LK et al., studied 170 hypertensive pregnant women to assess the efficacy of the dipstick method in detecting preeclampsia. The study concluded that the dipstick method has a significantly low PPV, with increased false-negative incidences [26]. A similar study by Jones C et al., concluded that the urine dipstick is not an adequate test for the diagnosis of proteinuria [27]. The present results further align with studies by White SL et al., and Siedner MJ et al., which reported that urine dipstick testing has poor sensitivity (57.8% and 63%, respectively) and relatively better specificity (95.4% and 86.1%, respectively), making it unreliable as a confirmatory diagnostic tool [28,29].

In comparison, UPCR showed significantly better sensitivity (91%) than the dipstick test (43%), although its specificity remained low at 7%. This aligns with a study by Chadha A and Tayade S which reported that UPCR has a sensitivity of 79.37%, a PPV of 92.59% and an accuracy of 75.79% [30]. A further study by Côté AM et al., which compared 13 studies in a systematic review, concluded that UPCR is an acceptable first-line diagnostic tool for identifying proteinuria, as 24-hour urine collections are frequently inaccurate, with discrepancies ranging from 13-68%, especially for those with a lean body weight. Among 161 women with serial 24-hour urine samples, 24.8% exhibited between-measurement differences of 25% or greater, exceeding both analytic and biological variation [31].

The present study findings are further supported by similar studies from Sanchez-Ramos L et al., which reported sensitivities and specificities for UPCR as 91.0% (95% CI 87.0-93.9) and 86.3% (95% CI 78.4-91.7), respectively [32]. Saudan PJ et al., also reported that UPCR had a positive correlation with 24-hour urine proteinuria (r=0.93, p<0.001) and demonstrated a sensitivity of 93% and specificity of 92% [33]. Other studies, such as that by Stefańska K et al., (sensitivity and specificity of 89% (95% CI, 75-97) and 100% (95% CI, 87-100), respectively) and Yamasmit W et al., (sensitivity, specificity and accuracy of 96.6%, 92.3% and 95.2%, respectively) also support these findings [34,35].

The F1-score, which balances sensitivity and precision, was notably higher for UPCR (59%) compared to the dipstick test (39%), reinforcing the notion that UPCR may be more useful for confirming proteinuria, although it may be less effective in detecting all true cases. These findings are in line with a study by Olisa CL et al., which demonstrated that UPCR has superior diagnostic accuracy compared with the dipstick method [19]. In a multicentre prospective study involving 2212 urine samples conducted by Baba Y et al., it was shown that the dipstick produced significantly higher false positive results compared to UPCR results [36]. The authors concluded that patients who show positive results on the dipstick should undergo further evaluation using UPCR results, even regardless of whether the patient is hypertensive or normotensive.

Limitation(s)

The present study has several limitations that should be considered when interpreting the data. The sample size of 80, being small, limits the ability to generalise the findings to a larger population. Additionally, while the 24-hour urine protein test is considered the gold standard, its accuracy remains a subject of debate [17]. The study was conducted at a single centre, underscoring the need for further multicentre studies to corroborate the findings.

CONCLUSION(S)

While the urine dipstick test and UPCR offer rapid alternatives, their diagnostic accuracy remains suboptimal when compared to the 24-hour urine protein test. Although UPCR is better at confirming proteinuria, it fails to reliably rule out preeclampsia due to its low NPV. These findings emphasise the need for a combination of clinical assessment, laboratory markers and the use of UPCR over the dipstick test as an adjunct.

REFERENCES

- [1] Karrar SA, Martingano DJ, Hong PL. Preeclampsia. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Aug 6]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK570611/.
- [2] Mathew R, Devanesan BP, Srijana, Sreedevi NS. Prevalence of hypertensive disorders of pregnancy, associated factors and pregnancy complications in a primigravida population. Gynecol Obstet Clin Med. 2023;3(2):119-23.
- [3] Malik A, Jee B, Gupta SK. Preeclampsia: Disease biology and burden, its management strategies with reference to India. Pregnancy Hypertens. 2019;15:23-31.
- [4] Shandilya V, Sinha N, Rani S. Preeclampsia: Prevalence, risk factors, and impact on mother and fetus. Indian J Cardiovasc Dis Women. 2023;8(3):193-99.
- [5] Battarbee AN, Sinkey RG, Harper LM, Oparil S, Tita ATN. Chronic hypertension in pregnancy. Am J Obstet Gynecol. 2020;222(6):532-41.
- [6] Eastern Virginia Medical School. Aspirin for the prevention of preeclampsia in women with stage 1 Hypertension: A pilot study [Internet]. clinicaltrials.gov; 2021 Jun [cited 2025 Feb 5]. Report No.: NCT04908982. Available from: https:// clinicaltrials.gov/study/NCT04908982.
- [7] Hofmeyr GJ, Belfort M. Proteinuria as a predictor of complications of preeclampsia. BMC Med. 2009;7:11.
- [8] Fishel Bartal M, Lindheimer MD, Sibai BM. Proteinuria during pregnancy: Definition, pathophysiology, methodology, and clinical significance. Am J Obstet Gynecol. 2022;226(2):S819-S834.
- [9] Müller-Deile J, Schiffer M. Preeclampsia from a renal point of view: Insides into disease models, biomarkers and therapy. World J Nephrol. 2014;3(4):169-81.
- [10] Haider MZ, Aslam A. Proteinuria. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 [cited 2025 Feb 4]. Available from: https://www.ncbi.nlm.nih.gov/books/NBK564390/.
- [11] Evans TRJ, Kudo M, Finn RS, Han KH, Cheng AL, Ikeda M, et al. Urine protein:creatinine ratio vs 24-hour urine protein for proteinuria management: Analysis from the phase 3 REFLECT study of lenvatinib vs sorafenib in hepatocellular carcinoma. Br J Cancer. 2019;121(3):218-21.
- [12] Olayinka L, Garnett E, Burnett B, Devaraj S. Comparison of random urine protein/ creatinine ratio with 24-hour urine protein in suspected pre-eclampsia. Pract Lab Med. 2023;36:e00316.
- [13] Gutiérrez-Peredo GB, Montaño-Castellón I, Gutiérrez-Peredo AJ, Lopes MB, Tapioca FPM, Guimaraes MGM, et al. The urine protein/creatinine ratio as a reliable indicator of 24-h urine protein excretion across different levels of renal function and proteinuria: The TUNARI prospective study. BMC Nephrol. 2024;25(1):418.
- [14] Clark WF, Macnab JJ, Sontrop JM, Jain AK, Moist L, Salvadori M, et al. Dipstick proteinuria as a screening strategy to identify rapid renal decline. J Am Soc Nephrol JASN. 2011;22(9):1729-36.
- [15] Lim D, Lee DY, Cho SH, Kim OZ, Cho SW, An SK, et al. Diagnostic accuracy of urine dipstick for proteinuria in older outpatients. Kidney Res Clin Pract. 2014;33(4):199-203.
- [16] Dimitriadis E, Rolnik DL, Zhou W, Estrada-Gutierrez G, Koga K, Francisco RPV, et al. Pre-eclampsia. Nat Rev Dis Primer. 2023;9(1):01-22.
- [17] Chang KJ, Seow KM, Chen KH. Preeclampsia: Recent advances in predicting, preventing, and managing the maternal and fetal life-threatening condition. Int J Environ Res Public Health. 2023;20(4):2994.
- [18] Osungbade KO, Ige OK. Public health perspectives of preeclampsia in developing countries: Implication for health system strengthening. J Pregnancy. 2011;2011;481095.
- [19] Olisa CL, Nwosu BO, Eleje GU, Oguejiofor CB, Mbachu II, Ogabido CA, et al. Comparison of urine protein-creatinine ratio and urine dipstick test for significant proteinuria in preeclamptic women. Ther Adv Reprod Health. 2024;18:26334941241288841.
- [20] Xiao J, Fan W, Zhu Q, Shi Z. Diagnosis of proteinuria using a random urine proteincreatinine ratio and its correlation with adverse outcomes in pregnancy with preeclampsia characterized by renal damage. J Clin Hypertens. 2022;24(5):652-59.
- [21] Ji S, Yang Y, Jeong Y, Hwang SH, Kim MC, Kim Y. Comparison of three types of analyzers for urine protein-to-creatinine ratios in dogs. J Vet Sci. 2021;22(1):e14.
- [22] Duckitt K, Harrington D. Risk factors for pre-eclampsia at antenatal booking: Systematic review of controlled studies. BMJ. 2005;330(7491):565.
- [23] Chan TF, Tung YC, Wang SH, Lee CH, Lin CL, Lu PY. Trends in the incidence of pre-eclampsia and eclampsia in Taiwan between 1998 and 2010. Taiwan J Obstet Gynecol. 2015;54(3):270-74.
- [24] Sheen JJ, Huang Y, Andrikopoulou M, Wright JD, Goffman D, D'Alton ME, et al. Maternal age and preeclampsia outcomes during delivery hospitalizations. Am J Perinatol. 2020;37(1):44-52.
- [25] Teeuw HM, Amoakoh HB, Ellis CA, Lindsley K, Browne JL. Diagnostic accuracy of urine dipstick tests for proteinuria in pregnant women suspected of preeclampsia: A systematic review and meta-analysis. Pregnancy Hypertens. 2022;27:123-30.

- [26] Phelan LK, Brown MA, Davis GK, Mangos G. A prospective study of the impact of automated dipstick urinalysis on the diagnosis of preeclampsia. Hypertens Pregnancy. 2004;23(2):135-42.
- Jones C, Jakubowski BE, Stevens R, Roberts N, McManus RJ, Tucker KL. The diagnostic accuracy of visual versus automated dipstick proteinuria testing in pregnancy: A systematic review and Meta-Analysis. Pregnancy Hypertens. 2024;35:73-81.
- White SL, Yu R, Craig JC, Polkinghorne KR, Atkins RC, Chadban SJ. Diagnostic accuracy of urine dipsticks for detection of albuminuria in the general community. Am J Kidney Dis Off J Natl Kidney Found. 2011;58(1):19-28.
- Siedner MJ, Gelber AC, Rovin BH, McKinley AM, Christopher-Stine L, Astor B, et al. Diagnostic accuracy study of urine dipstick in relation to 24-hour measurement as a screening tool for proteinuria in lupus nephritis. J Rheumatol. 2008;35(1):84-90.
- Chadha A, Tayade S. Urinary protein-to-creatinine ratio: An indicator of adverse clinical outcomes in preeclampsia with proteinuria. Cureus. 2022;14(3):e23341.
- Côté AM, Firoz T, Mattman A, Lam EM, von Dadelszen P, Magee LA. The 24hour urine collection: Gold standard or historical practice? Am J Obstet Gynecol. 2008;199(6):625.e1-6.

- [32] Sanchez-Ramos L, Gillen G, Zamora J, Stenyakina A, Kaunitz AM. The proteinto-creatinine ratio for the prediction of significant proteinuria in patients at risk for preeclampsia: A meta-analysis. Ann Clin Lab Sci. 2013;43(2):211-20.
- Saudan PJ, Brown MA, Farrell T, Shaw L. Improved methods of assessing proteinuria in hypertensive pregnancy. Br J Obstet Gynaecol. 1997;104(10):1159-64.
- [34] Stefańska K, Zieliński M, Zamkowska D, Adamski P, Jassem-Bobowicz J, Piekarska K, et al. Comparisons of dipstick test, urine protein-to-creatine ratio, and total protein measurement for the diagnosis of preeclampsia. Int J Environ Res Public Health. 2020;17(12):4195.
- Yamasmit W, Chaithongwongwatthana S, Charoenvidhya D, Uerpairojkit B, Tolosa J. Random urinary protein-to-creatinine ratio for prediction of significant proteinuria in women with preeclampsia. J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet. 2004;16(5):275-79.
- Baba Y, Yamada T, Obata-Yasuoka M, Yasuda S, Ohno Y, Kawabata K, et al. Urinary protein-to-creatinine ratio in pregnant women after dipstick testing: Prospective observational study. BMC Pregnancy Childbirth. 2015;15:331.

PARTICULARS OF CONTRIBUTORS:

- Assistant Professor, Department of Obstetrics and Gynaecology, Dr. D. Y. Patil Medical College, Dr. D. Y. Patil Vidyapeeth (Deemed to be University), Pimpri, Pune, Maharashtra, India.
- Professor, Department of Obstetrics and Gynaecology, Dr. D. Y. Patil Medical College, Dr. D. Y. Patil Vidyapeeth (Deemed to be University), Pimpri, Pune, Maharashtra, India.
- Resident, Department of Obstetrics and Gynaecology, Dr. D. Y. Patil Medical College, Dr. D. Y. Patil Vidyapeeth (Deemed to be University), Pimpri, Pune, Maharashtra, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Pragya Tiwari,

Resident, Department of Obstetrics and Gynaecology, Dr. D. Y. Patil Medical College, Hospital and Research Centre, Dr. D. Y. Patil Vidyapeeth (Deemed to be University), Pimpri, Pune-411018, Maharashtra, India.

E-mail: pragyatiwariajoy@gmail.com

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. NA

PLAGIARISM CHECKING METHODS: [Jain H et al.] ETYMOLOGY: Author Origin

- Plagiarism X-checker: Feb 05, 2025
- Manual Googling: Mar 19, 2025
- iThenticate Software: Mar 21, 2025 (11%)

EMENDATIONS: 6

Date of Submission: Feb 04, 2025 Date of Peer Review: Feb 22, 2025 Date of Acceptance: Mar 23, 2025 Date of Publishing: Apr 01, 2025