

Umbilical Cord Coiling Index and Perinatal Outcome

NIVEDITA S. PATIL¹, SUNANDA R. KULKARNI², RENU LOHITASHWA³

ABSTRACT

Objectives: To evaluate the perinatal outcome with the abnormal umbilical cord coiling index.

Study Design: This prospective study was carried out in the department of OBG at Adichunchangiri Institute of Medical Sciences, B.G.Nagara, Mandya, Karnataka, India from January 2008 to August 2010. 200 patients who were in active labour with term gestations, irrespective of their parities, who had singleton pregnancies with live babies who were either delivered by vaginal or LSCS were included in the study. Umbilical cord coiling index was calculated and it was correlated with various perinatal parameters like birth weight, meconium stained liquor, Apgar score, ponderal index and foetal growth restriction. Chi square and Fisher exact tests were used to find the significance

of study parameters.

Results: There was a significant correlation between the hypercoiled cords (UCI >90th percentile) and IUGR of the babies (p value of < 0.001) and low ponderal indices (a p value of 0.022) Hypocoiled cords (UCI which was < 10th percentile) were significantly associated with meconium staining (p < 0.001), Apgar score at 1 min of <4 and at 5 min of <7 (p value 0.065), LSCS rates (p value of 0.008) and NICU admissions (p <0.001).

Conclusion: Hypercoiled cords or UCI which was > 90th percentile was associated with IUGR and low ponderal indices. Hypocoiled cords or UCI which was <10th percentile was associated with meconium staining, Apgar score at 1 min of <4 and at 5 min of <7, more LSCS rates and more NICU admissions.

Key words: Umbilical cord coiling index, Hypercoiling umbilical cord, Hypocoiling umbilical cord

INTRODUCTION

The umbilical cord is vital for the development, wellbeing and survival of the foetus and yet, it is vulnerable to kinking, compressions, traction and torsion, which may affect the perinatal outcome. vessels are the most vulnerable part of the foetal anatomy. The total number of coils for any particular cord is believed to be established early in gestation. The pattern of coiling develops during second and third trimesters, presumably due to snashes in the cord and this coiling changes as pregnancy advances. The three blood vessels pass along the length of the cord in helical or coiled fashions [1]. The helical fusion of these umbilical vessels is termed as spiral course [2].

A coil is defined as having completed a 360 spiral course of umbilical vessel around Wharton's jelly. Coiling property of umbilical cord was described by Berengarius in 1521 [1]. In 1954, umbilical coiling was first quantified by Edmonds who divided the total number of coils by umbilical cord length in centimetres and called it "Index of twist". He assigned positive and negative scores to clockwise and anticlockwise coiling, respectively [1,3]. Later, Strong et al simplified it by eliminating three directional score and named it "The umbilical cord coiling index" [1,4]. An abnormal umbilical cord coiling index includes both hypocoiled cords (cords with an umbilical cord coiling index which is < 10th percentile) and hypercoiled cords (cords with an umbilical cord coiling index which is > 90th percentile). An abnormal umbilical coiling index has been reported to be related to adverse perinatal outcomes [1,2,5-7].

The present study aimed to evaluate the perinatal outcomes with abnormal umbilical cord coiling indices.

MATERIAL AND METHODS

This was a prospective study which was carried out in the Department of Obstetrics & Gynaecology at Adichunchangiri Institute of Medical Sciences, B.G.Nagara, Mandya, Karnataka, India, from January 2008 to August 2010. Ethical committee

clearance was obtained from the Institute's Research Council and Ethics Committee. 200 patients who were in active labour with term gestations, irrespective of their parities, who had singleton pregnancies with live babies who were either delivered by vaginal or LSCS were included in the study. Patients were observed in second and third stages of labour.

After separating baby from umbilical cord, the cord was tied and cut as close to baby as possible. The umbilical cord was measured in its entirety, including the length of the placental end of the cord and umbilical stump on the baby. Number of complete coils or spirals were counted from neonatal end towards the placental end of cord and it was expressed in cm. After this, UCI was calculated by dividing the total number of coils by the total length of cord in cm.

$$\text{UCI} = \frac{\text{Number of coils}}{\text{Total length of cord in cms}}$$

Then, perinatal factors like meconium staining, foetal weight, Apgar score and ponderal index were correlated with UCI. Thus, the effect of umbilical vascular coiling on perinatal outcome was carried out. All patients and babies were followed till they were discharged.

RESULTS

Among 200 women who were studied, 137 were booked and 63 were unbooked. Primigravida were 129 and multigravida were 71. 110 delivered vaginally, 79 delivered by LSCS, 10 had vacuum deliveries and 1 had an outlet forceps delivery. 90 women who had undergone either LSCS or instrumental deliveries had indications of foetal distress.

In the present study, UCI which was < 10th percentile < 0.06, was considered as hypocoiling whereas UCI which was > 90th percentile

> 0.48 was considered as hypercoiling. 156 cords showed normal coiling, 23 showed hypocoiling, and 21 showed hypercoiling. 22 babies weighed < 2.5kg. Meconium staining liquor was present in 42 cases. An Apgar score at 1 min of < 4 was observed in 31 cases and that of > 4 was seen in 169 cases. Apgar score at 5 min was < 7 in 28 cases and it was > 7 in 172 cases. The twist of the cord was dextral in 77 cases and it was sinistral in 123 cases. 35 babies were admitted to the NICU. 108 were males and 92 were females.

[Table/Fig-1] (in row A,B,C,D) show correlation of meconium staining, Apgar scores at 1 min, Apgar scores at 5 min and NICU admissions with UCI respectively.

This being statistically significant, it suggests that hypocoiled cords are more associated with meconium staining, low Apgar scores and more NICU admissions of babies. [Table/Fig-1] (in row E and F) show correlation of IUGR and ponderal indices with UCI respectively. These are statistically significant, suggesting that hypercoiled cords are associated with IUGR and low ponderal indices.

[Table/Fig-2] shows correlation of UCI with mode of delivery. Hypocoiled cords were significantly associated with incidence of more LSCS.

Baby details	UCI			P value
	Normal	Hypo coiling	Hyper coiling	
A. Meconium staining				
Yes	17(10.89%)	23(100%)	2(9.5%)	<0.001
No	139(89.11%)	0(0%)	19(90.5%)	
B. Apgar score at 1 min				
<4	23(14.7%)	7(30.4%)	1(4.8%)	0.072
>4	133(85.3%)	16(69.6%)	20(95.2%)	
C. Apgar score at 5 min				
<7	19(12.2%)	7(30.4%)	2(9.5%)	0.065
>7	137(87.8%)	16(69.6%)	19(90.5%)	
D. NICU admission				
Yes	19(12.2%)	10(43.5%)	6(28.6%)	<0.001
No	137(87.8%)	13(56.5%)	15(71.4%)	
E. IUGR				
Yes	5(3.2%)	0(0%)	17(81%)	<0.001
No	151(96.8%)	23(100%)	4(19%)	
F. Ponderal index				
<2.5	108(69.2%)	15(65.2%)	20(95.2%)	0.022
>2.5	48(30.8%)	8(34.8%)	1(4.8%)	

[Table/Fig-1]: Correlation of Meconium Staining, Apgar score, NICU admission, IUGR and Ponderal index with UCI

Maternal variables	UCI			P value
	Normal	Hypo coiling	Hyper coiling	
Mode of delivery				
FTVD/ FTVMD/ FTVOD	101(64.7%)	7(30.4%)	13(61.9%)	0.008**
LSCS	55(35.3%)	16(69.6%)	8(38.1%)	

[Table/Fig-2]: Correlation of mode of delivery with UCI

DISCUSSION

Several trials in the past have correlated the relationship between perinatal outcomes and the UCI. In the present study, UCI was compared with various parameters. On comparing UCI with parity, it was found that there was no statistical significance between primigravida and multigravida and that there was no statistical significance between the dextral and sinistral twists of cord. In the present study, hypocoiled cords were significantly associated

with LSCS. In the present study, it was observed that meconium staining was significantly associated with UCI which was <10th percentile. Gupta S et al., studied 107 umbilical cords and found that in hypocoiled group, meconium staining was significantly higher than in those with normocoiled group [1]. Strong TH et al., studied 100 cases and they found that meconium staining was associated with UCI values of less than 10th percentile, with a p value of 0.03, which was highly significant [4]. In another study which was done by Padmanabhan LD et al., 130 cases were studied, where they found that meconium staining was significant in hypercoiled group [8].

In the present study, Apgar score at 1 min of < 4 was found with UCI which was < 10th percentile. Gupta S et al., studied 107 umbilical cords and found that in hypocoiled cords, low Apgar scores were present [1]. In another study which was done by Padmanabhan et al., 130 umbilical cords were studied and it was found that in hypocoiled group, there were significantly low Apgar scores [8].

In the present study, Apgar score at 5 min of < 7 in relation to UCI was seen with < 10th percentile. Monique WM et al., studied 885 cases and found that hypocoiling was associated with low Apgar scores [8,9]. Gupta S et al., studied 107 cords and found that babies with Apgar scores of < 7 had significantly lower UCIs than the babies with Apgar scores of > 7 [1]. Padmanabhan LD et al., studied 130 umbilical cords and found that hypocoiled group was associated with low Apgar scores of < 7 [8].

In the present study, it was observed that IUGR was significantly associated with UCI which was >90th percentile. In a similar study which was done by Monique WM et al., 885 patients were studied and it was found that hypercoiling was associated with small for gestational age infants [8/9]. Georgiou AM et al., studied 34 cords and it was found that hypercoiled cords were associated with IUGR [10].

In the present study, babies who were admitted to NICU were significantly associated with UCI which was < 10th percentile. Monique WM et al., studied 885 cases and it was concluded that hypocoiling of the cord was associated with NICU admissions of foetal death [8]. Strong TH et al., studied 687 cases and it was found that incidence of foetal death in non coiled group was significantly greater, with a p value of <0.05 [11].

In the present study, cords with UCI which was >90th percentile had low ponderal indices. As ponderal index is low in foetal growth restriction, hypercoiled cords are associated with both foetal growth restriction and low ponderal indices. Gupta S et al., studied 107 cords and it was concluded that there was no statistically significant difference between the hypocoiled group and hypercoiled group in association with ponderal indices [1].

CONCLUSION

The abnormal umbilical coiling index is associated with adverse perinatal outcomes. UCI which is > 90th percentile is termed as hypercoiling and it is associated with foetal growth restriction. It was found that a ponderal index of < 2.5 was associated with hypercoiling of cord. UCI which is < 10th percentile is termed as hypocoiling and it is associated with meconium staining, low Apgar scores and NICU admissions of babies. Therefore, an antenatal detection of coiling index can identify fetuses who are at risk and thus help in further management.

REFERENCES

- [1] Gupta S., Faridi MMA, Krishnan J., Umbilical Coiling Index. *J. Obstet Gynecol India*. 2006; 56 (4) : 315-19.
- [2] Rana J, Ebert GA, Kappy KA. Adverse perinatal outcome in patients with an abnormal umbilical coiling index. *Obst. Gynaecol*. 1995 ; 85 (4); 573-77.
- [3] Edmonds HW. The spiral twist of normal umbilical cords in twins and singlestone. *Am. J. Obstet Gynecol*. 1954 ; 67 : 102-20.
- [4] Strong TH, Jarles DL, Vega JS Feldman DB. The umbilical coiling index. *Am. J. Obst. Gynecol Part I*. 1994 ; 170 (1) : 29-32.

- [5] Larco RV, Jones KL, Benirschke K. The umbilical cord twist : origin, directions, and relevance. *Am. J. Obstet Gynaecol.* Part I 1987; 157 (4) : 833-38.
- [6] Strong TH, Finberg HL, Mattox JH. Antepartum diagnosis of noncoiled umbilical cords. *Am J. Obstet Gynecol.* 1994 ; 170 : 1729-33.
- [7] Ercal T, Lacin S, Altungyurt S et al., Umbilical coiling index : is it marked for foetus at risk? *Br J Clin Pract.* 1996 ; 50 : 254-56.
- [8] Pudmanabhan LD, Mhaskar R, Mhaskar A. Umbilical vascular coiling and the perinatal outcome. *J Obstet Gynecol India.* 2001; 51 (6): 43-44.
- [9] Monique WM, de Laat, Frank A, Dots ML, Verses GHA, Nikkels GJN. *Obstet Gynecol.* 2006; 107 (5) : 1049-55.
- [10] Georgiou HM, Rice GE, Walker SP, Wein P, Neil M, Permezel M. The effect of vascular coiling on venous perfusion during experimental umbilical cord encirclement. *Am. J. Obstet Gynecol.* 2001; 184 (4): 673-78.
- [11] Strong TH, Elliott JP, Radin TG. Non-coiled umbilical blood vessels: A new marked for fetus at risk. *Obstet Gynecol.* 1993 ; 81 : 409-11.

PARTICULARS OF CONTRIBUTORS:

1. Senior Resident, Department of Obstetrics & Gynaecology, SNMC, Bagalkot, Karnataka, India.
2. Professor, Department of Obstetrics & Gynaecology, AIIMS, B.G. Nagara, Mandya, Karnataka, India.
3. Assistant Professor, Department of Physiology, SSIMS&RC, Davangere, Karnataka, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Nivedita S. Patil,
Senior Resident, Department of Obstetrics & Gynaecology, SNMC, Bagalkot, India.
Phone: 9591492320, E-mail: nivvi_patil@yahoo.com

Date of Submission: **Oct 12, 2012**
Date of Peer Review: **Jan 28, 2013**
Date of Acceptance: **Jun 17, 2013**
Date of Publishing: **Aug 01, 2013**

FINANCIAL OR OTHER COMPETING INTERESTS: None.