

Evaluation of Antibiotic Residues in Pasteurized and Raw Milk Distributed in the South of Khorasan-e Razavi Province, Iran

MORTEZ MOHAMMADZADEH MOGHADAM¹, MOSTAFA AMIRI²,
HAMED RAMEZANI AWAL RIABI³, HAMID RAMEZANI AWAL RIABI⁴

ABSTRACT

Introduction: The presence of antibiotic residues in milk and other products livestock is a health problem which can endanger public health. Antibiotics are used widely in animal husbandry to treat diseases related to bacterial infections. Antimicrobial drugs have been in use for decades in industry. They are commonly used in livestock facilities to treat mastitis.

Aim: This study aimed to investigate antibiotic residues in pasteurized milk distributed in schools, in milk collection centers, and in milk production factories in Gonabad city.

Materials and Methods: This cross-sectional study was conducted on 251 samples of commercial pasteurized milk packet distributed in schools (code A), raw milk collection centers in Gonabad city (code B), and pasteurized milk production factories (code C) in Gonabad city. The Copan test kit of Denmark Christian Hansen Company was used to monitor antibiotic residues in milk. The data were analysed employing Chi-square test and one-way analysis of variance (ANOVA) to determine significant differences using SPSS software version 20. The significant level was considered at $p < 0.05$.

Results: In total, 251 milk samples were collected out of which 143 (57%) were code A, 84 (33.5%) code B and 24 (9.6%) code C. Total number of 189 samples (75.2%) were negative and 62 (24.8%) were positive. From the three types of milk samples, 41 samples (28.7%) of the code A, 18 samples (21.4%) of the code B and 3 samples (12.5%) of the code C were positive.

In general, from the milk samples most contaminated with antibiotics, 17 samples were positive in January and regarding code A, 13 samples were positive in the same month. There was not a significant difference among the three types of milk ($p > 0.05$). The highest number of milk samples ($n = 7$) contaminated with antibiotics were related to code B (38.5%). Most positive cases were related to code A in winter. Also, there was no significant difference among the three types of contaminated milk regarding the year and month ($p = 0.164$ and $p = 0.917$, respectively).

Conclusion: Pasteurized milk supplied in the studied city has high level of contamination due to high use of antibiotics. A standard limit needs to be set for the right level of residue of antibiotics in milk to avoid the harmful effects.

Keywords: Antibiotic residue, Copan test, Food safety, Milk quality

INTRODUCTION

Antibiotics are used widely in animal husbandry for treating the diseases related to bacterial infections [1,2]. Antimicrobial drugs have been used extensively for decades in industry. They are commonly used in livestock facilities to treat mastitis [3,4]. Annually around the world, about 12 million kg of antibiotics are used in the livestock industry, out of which only about 9 million kg are used to treat infectious diseases and the rest are used either to prevent or increase the animal growth [5].

According to statistics, 806 tons of antibiotics were used in animal husbandry and poultry in Iran in 2011, of which 66.4% were used in the livestock industry [6]. The presence of antibiotic residues in milk is a public health problem which can lead to allergic reactions in sensitive individuals and has negative effects on the human gut flora composition [3]. It has also been reported that tetracycline residues, in the long run, cause digestive disorders, the destruction of the liver, yellowing of the teeth and the tooth hyperplasia [7].

Antibiotic residues and veterinary drugs can trigger cancer, mutagenicity and toxicity in humans [8]. Further, the frequent use of antibiotics can increase the biological drug resistance [1,3]. Besides, antibiotic residues in animal products will cause environmental pollution and water pollution [9]. In most countries of the world, for the remainder of antimicrobial drugs in milk, some criteria have been set [1].

Due to the harmful effects of veterinary drugs for humans, the Food and Drug Administration System in Iran enacts laws to control the

use of antibiotics. Milk production in Iran is about 7 million tons and the consumption of edible milk per capita has reached to 4.82kg in recent years [5]. Currently, both raw and pasteurized milk should be free of microbial growth inhibitor according to Iran's internal standards set by the Iranian Food and Drug Administration System; however, a threshold for the remaining antimicrobials has not been set exclusively [10,11].


Due to the harmful effects of residual antimicrobial drugs in food especially in domestic animals' milk which is consumed with breakfast by many people, particularly by school children in Iran, quality control regarding the antibiotic residue in milk is essential. This study was designed to determine the level of antibiotic residues in milk and increase monitoring of producers of dairy products through systematic planning so that the health of milk consumers, who are mainly school students, could be protected.

MATERIALS AND METHODS

This study is a cross-sectional one; the sampling was performed according to the standard procedures set by the Institute of Standards and Industrial Research of Iran-2836. Random cluster sampling method was used, and the sample size, based on prior studies [12], was calculated for every year by the following formula:

$$n = \frac{Z_{1-\alpha/2} P(1-P)}{d^2}$$

$$p = 21\%, 1-p = \%20, d = 0.06, Z = 1.96, n = 45$$



[Table/Fig-1]: Copan Kit reading guide (Forouzan, 2014) [13].

Total 251 samples of commercial pasteurized milk packet distributed in schools (code A), raw milk collection centers in Gonabad city (code B) and pasteurized milk production factories (code C) in Gonabad city were included in the study. The samples were collected in 2009-2013 years, and once a month, 200 micro liters of milk were collected randomly in sterile containers on which the product specifications were listed. The samples were divided into three groups, for which the brand name of the products were not mentioned to observe the ethical considerations. The milk samples were transferred to the laboratory at Food and Drug Administration Deputy. The Copan test kit of Denmark Christian Hansen Company was utilised to detect the presence of antibiotic residues in milk samples. Copan test, is a quick, particular, and applicable method, which is useful in regulatory and production centers for food products and is used to indicate the presence of the antibiotics by inhibiting microbial growth [13].

Aliquots of 100µL of milk samples were poured into the kit containing *Bacillus stearothermophilus* var. *calidolactis* using a micro-pipette, the kits were incubated at 64.5°C and the results were recorded after 3 hours. The residues of microbial growth were identified in the absence of antibiotics. During growth, the microorganism uses nutrients, ferments lactose and produces lactic acid which results in a change in the colour of the compound bromocresol from purple to yellow.

On the contrary, if the antibiotic residues exist, *Bacillus stearothermophilus* var. *calidolactis* unable to grow and the purple colour of the medium remains unchanged. The colour of suspected samples would be between purple and yellowish colour; hence, determining the presence of antibiotics type in these samples was not conclusive. The kit reading guide is specified in [Table/Fig-1] [13].

STATISTICAL ANALYSIS

Statistical analysis was carried out using Chi-square and one-way ANOVA to determine if there were any significant differences among the groups with SPSS software V.20. The significant level was considered at $p < 0.05$.


RESULTS

In general, 251 milk samples were collected, of which 143 (57%) were coded as A, 84 (33.5%) coded as B, and 24 (9.6%) coded as C. Of these, 189 samples (75.2%) were negative and 62 others (24.8%) were positive. From the three types of milk samples, 41 samples (28.7%) of the code A, 18 samples (21.4%) of the code B, and 3 samples (12.5%) of the code C were positive.

In addition, there was no significant difference among the three types of contaminated milk with regard to the year and month ($p = 0.164$ and $p = 0.917$, respectively) [Table/Fig-2]. Totally, from the milk samples most contaminated with antibiotics, 17/62 (27.4%)

Code sample	%Frequency	% Positive samples in year					p-value
		2009	2010	2011	2012	2013	
A	57	17.1	7.3	26.8	14.6	34.1	0.365
B	33.5	11.1	11.1	33.3	5.6	38.9	0.932
C	9.6	33.3	33.3	33.3	0	0	0.770


[Table/Fig-2]: Comparison of the positive samples in percent in the years (2009-13).



[Table/Fig-3]: The frequency of positive milk samples per month- year (2009-13).


Variable		% Positive samples					
		A	p-value	B	p-value	C	p-value
Season	Spring	7.3	0.628	16.6	0.725	33.3	0.311
	Summer	7.3		16.6		0	
	Autumn	19.5		22.3		33.3	
	Winter	65.9		44.5		33.4	
Month	Apr	2.4	0.736	0	0.673	0	0.676
	May	2.4		5.5		33.4	
	Jun	2.4		11.1		0	
	Jul	2.4		0		0	
	Aug	4.8		16.6		0	
	Oct	0		0		0	
	Sep	0		5.5		0	
	Nov	4.8		5.5		33.3	
	Dec	0		11.1		0	
	Jan	31.7		16.6		33.3	
	Feb	14.6		16.6		0	
	Mar	29.2		11.1		0	

[Table/Fig-4]: Comparison of the percentage of positive milk samples collected in season and month.



[Table/Fig-5]: The frequency of positive and negative samples of milk per year.

Neg - Negative
Pos - Positive



[Table/Fig-6]: The frequency of positive and negative milk samples according to season.

Neg - Negative
Pos - Positive

samples were positive in January, and regarding code A, 13 samples were positive in the same month [Table/Fig-3].

There was not a significant difference among the three types of milk ($p>0.05$). The highest number of milk samples ($n=7$) contaminated with antibiotics were related to code B (38.5%). During winter, the highest number of positive milk samples belonged to code A and B, 27 (65.9%) and 8 (44.5%) samples, respectively [Table/Fig-4].

Considered as a whole, the highest number of milk samples contaminated with antibiotics were in the year 2013 ($n=21$ samples, 33.8%). The proportion of positive samples to the whole samples in the years (2009, 2010, 2011, 2012 and 2013) were 10 (28.7%), 6 (26.9%), 18 (20.9%), 7 (19.3%) and 21 (28.5%), respectively [Table/Fig-5].

The proportion of positive milk samples regarding the entire sample in the different seasons of the year were 36 (26.4%), 13 (25.4%), 7 (24.1%) and 6 (17.1%) in winter, autumn, spring, and summer, respectively [Table/Fig-6].

DISCUSSION

Milk is a perfect food, which provides the bulk of the daily requirements for humans of all ages, especially children. On the other hand, not only milk is a good source for the growth of microorganisms but it is also a source of food poisoning for health, so its importance is quite clear [14]. According to international standards, the presence of antibiotics in both milk and milk products is not acceptable because consumers may already have allergic reactions and gastrointestinal problems. Upon consumption by livestock, almost all antibiotics appear in the raw milk in shortest period of time. The presence of antibiotics in milk can upset the process of dairy production. Therefore, the presence of antibiotics in the raw milk should be controlled by the factories producing pasteurized milk.

The results of this study showed that over five years, 23% of pasteurized milk samples in Gonabad city were positive regarding antibiotics. Between the months of the year in terms of milk samples contaminated with antibiotics, there was no significant difference; hence, it seems that climate change has no impact on the results. The milk coded as A, that is, the one included the milk distributed in schools, had the highest contamination that is not enjoyable for students and makes them refrain from consuming milk. On the

whole, there was no significant difference among samples over the years in terms of contamination of milk with antibiotics, a finding which indicates that there is not enough control over the use of antibiotics in veterinary medicine. In other words, the degree of consuming antibiotics was expected to be low with the course of time; however, it was not observed in our studied results.

The literature shows that most of the research has investigated the raw milk in addition to the pasteurized one. Furthermore, in the previous studies, mostly, the sampling method has been different from the one in our study and the milk consumed by the school children has not been investigated. The results of the present study are in some cases similar to those of other studies and in some other cases the percent of positive samples has been low [15]. In what follows, some of these studies are mentioned. The results of our study were compared with those of other studies in Iran [Table/Fig-7] [16-21].

Lee Burton et al., reported that the two tests of Delvo and Copan can detect milk contamination to Penicillin, Cloxacillin Sodium, Sulfamethazine, Cephalexin and Gentamycin based on the standards set by the European Union. These two tests are simple, low-error, and are complementary for the needs of the European Union [22]. In the study Kangth et al., in Kenya, it was shown that 16% of milk samples contained antibiotic residues [7].

Bylandzic et al., reported that, in 1259 samples of raw milk in Croatia, 37 cases of residual antibiotics was diagnosed [23], nevertheless, the raw milk was not examined in this study.

The results obtained in a study by Mahmoudi et al., also, indicated that significant contamination with antibiotic residues in pasteurized milk samples was observed, that is, 22.2% of the samples were positive in terms of contamination [12]. In another study in Iran, from a total of 848 milk samples, 30% cases were eligible for antibiotic residues.

Movassegh (2012) tested 50 samples of raw cow milk in the Ilkhchi region in Tabriz using Copan test and observed that 5 samples (10%) contained antibiotic residues [8]. Rasouli et al., reviewed the status of antibacterial drugs on dairies in Tehran and reported that among 24 anti-bacterial products, Tetracycline, Penicillin, and Streptomycin, respectively, had the highest frequency of use. Most

Researcher	Reference number	Year	Sample type	Test	Milk contamination (%)	Compared with our study
Present	-	2009-2013	251 Samples of raw and pasteurized milk in Gonabad city	Copan	28.7% of schools, 21.4% of collection centers, 12.5% of factories	-
Mahmoudi	[16]	2012	200 samples of raw milk dairies in the outskirts of the Tabriz city	Delvo test	57.5%	Different method and results
Mahmoudi	[17]	2011	Samples of raw and pasteurized milk in Ilam Province	Copan	29.1% of milk in collection centers and 22.2% of pasteurized milk	Antibiotic residues of milk was higher in summer than that of the spring (similar to our study)
Manafi	[18]	2008	Raw and pasteurized milk East Azerbaijan Province	Delvo test	26% of industrial dairies, 16% of collection centers, and 30% of pasteurized milk	In our study, the percentage of contaminated milk samples was more than that of this study
Bahreinpour	[19]	2008	200 samples of raw and pasteurized milk collected in Mashhad city	Microbial inhibition test	30%	Different method and results
Ghanavi	[20]	2003	Milk samples collected in Iran	Different methods	27% of raw and 5% of pasteurized milk	Different method and results
Abedini	[21]	1973	800 samples of raw and pasteurized milk in Shiraz city	Disc method	2.7%	The samples were collected in 8 months (in our study during 12 months)

[Table/Fig-7]: Comparison of the results of our study with those of other studies in Iran [16-21].

of these husbandry centers do not follow the right procedure and time for treatment with antibiotics and the proper time interval for consumption or prohibition of both milk products and meat [24,25].

The results of Fallahrad et al., conducted on 68 milk samples, pointed out that 11.76% of the total samples of the raw and pasteurized milk collected in Mashhad city were contaminated with Gentamycin [26]. In a study by Dabbagh Moghaddam et al., (2014), which was performed on 56 samples of pasteurized milk distributed in Tehran, the researchers reported the presence of antibiotics in 33.9% of the sample, of which 26.3% of the samples were contaminated with Tetracycline and 7.6% of the samples contained other antibiotics [27]. Sadeghi et al., reported that, from a total of 100 samples of raw milk taken from the milk tankers in Garmsar, 19% of the samples were contaminated with residues of antibiotics [28]. The results of the study carried out by Mohammadi Sani et al., showed that 40.8% of pasteurized milk samples, which were tested with Copan test, were positive [29]. In Yamaky's study which was performed on 2686 samples of raw sheep's milk in Spain, only 7.1% of the samples were found to contain residues of antibiotics [30].

The results of the above studies conducted in Iran shows that, at present, antimicrobial residues in milk production is high and the results of our study confirm these findings. Therefore, the prevalence of antibiotic residues in raw and pasteurized milk production in Iran is striking and disturbing.

The results of studies in other countries, especially European countries indicate that controlling and monitoring milk production in these countries are more serious, and the dosage and timing of the antibiotics are observed more precisely. Given that milk a complete food and that the nutritionists insist that it should be used every day by people of all ages, especially children, the pasteurized milk is distributed free of charge in Iranian schools. However, the control of antimicrobial residues in milk is an important and necessary issue.

Currently, manufacturers of dairy products usually use high quality milk for the production of other milk products because the antimicrobial residues can inhibit the growth of starters or the useful bacteria needed for producing those products [31].

LIMITATION

One limitation of this study was that there were some undiagnosed antibiotics which were not specified in livestock products due to lack of access to appropriate diagnostic tests. Also, another limitation was that we did not include unpasteurized raw milk in our sample as it was not available and the milk production centers were so dispersed.

CONCLUSION

Finding of above study noted that the production of contaminated pasteurized milk in Gonabad city is above the standard level, which is due to the high use of antibiotics in livestock. There is not sufficient and necessary control over the milk factories and milk collection centers in determining the remaining antibiotics during delivery of milk. On the other hand, poor livestock management, low supervision over the livestock and husbandries, non-compliance with sanitary conditions, and inadequate physical space can cause the animals to get diseases, for the treatment of which the farmers have no choice other than taking antibiotics.

Thus, providing necessary training for farmers and ranchers of livestock, supervising and controlling the construction of physical space for livestock by the relevant organizations, and also preventing the delivery of contaminated milk to the factories for milk products would be very effective in reducing antimicrobial residues in pasteurized milk. Moreover, control and detection of antibiotics in food and other products of animal origin (meat, poultry, fish, eggs, etc.), can also be done to protect the health of consumers. It is recommended that the right residue of any kind of antibiotics in Iran's internal standards be set similar to respective standards of the European Union.

ACKNOWLEDGMENTS

We should thank the staff at food laboratory of Food and Drug Department of Gonabad University of Medical Sciences for helping us to perform the tests.

REFERENCES

- [1] Stead SL, Ashwin H, Richmond SF, Sharman M, Langeveld PC, Barendse JP, et al. Evaluation and validation according to international standards of the Delvotests SP-NT screening assay for antimicrobial drugs in milk. *Inter Dai J*. 2008;18:3-11.
- [2] Zhang X, Chen L, Xu Y, Wang H, Zeng Q, Zhao Q, et al. Determination of B-lactam antibiotics in milk based on magnetic molecularly imprinted polymer extraction coupled with liquid chromatography-tandem mass spectrometry. *J of Chr B*. 2010;878:3421-26.
- [3] Chung H, Lee J, Chung Y, Lee K. Analysis of sulfonamide and quinolone antibiotic residues in Korean milk using microbial assays and high performance liquid chromatography. *J F che*. 2009;113:297-301.
- [4] Pengov A, Kirbis A. Risks of antibiotic residues in milk following intramammary and intramuscular treatments in dairy sheep, *analytica chimica acta*, 2009; 637:13-17.
- [5] Karim G, kiaei SMM, Rokni N, Razavi Rouhani SM. Milk contaminated with residues of antibiotics during the last forty years in Iran. *J Heal Food*. 2011;1(1):23-30.
- [6] Aalipour F, Mirlohi M, Jalali M. Prevalence of antibiotic residues in commercial milk and its variation by season and thermal processing methods. *Int J of Env Hea Eng*. 2013;2(1):41.

- [7] Kang'ethe EK, Aboge GO, Arimi SM, Kanja LW, Omoro AO, McDermott JJ. Investigation of the risk of consuming marketed milk with antimicrobial residues in Kenya. *Food Control*. 2005;16:349-55.
- [8] Movasseh MH. Detection of antibiotic residue in raw cow milk in Ilikhchi (South West Tabriz) in the spring of 2009. *J of F Tech & Nut*. 2012;9(3):89-94. [In Persian].
- [9] Kemper N, Farber H, Skutlarek D, Krieter J. Analysis of antibiotic residues in liquid manure and leachate of dairy farms in Northern Germany. *Agri Wat Man*. 2008;95:1288-92.
- [10] Institute of standards and industrial research of Iran, milk and milk products – raw milk – specification and test methods, no: 164:1383
- [11] Institute of standards and industrial research of Iran, pasteurized milk specifications and test methods, no: 93; 1386.
- [12] Mahmoudi R, Amini K, Vahabzade M, Mir H, Vagef R. Antibiotic residues in raw and pasteurized milk, Iran. *J of Res & Health*. 2014;4(4):884-89.
- [13] Forouzan S, Rahimirad A, Seyedkhoei R, Asadzadeh J, Bahmani M. Determination of antibiotic residues in the pasteurized milk produced in. *J of Coa Lif Med*. 2014;2(4):297-301.
- [14] Zolfaghari MR, Gaeini R, Kalhor N, Khalilian, Razavian MH. Study of microbial contamination of milk and pasteurization dairy products in Qom province. 2012;5(1,2):47-57.
- [15] Mahmoudi R, Golchin A, Farhoodi A. A review on antibiotic residues in animal-derived Foods in Iran over the last thirty years. *Journal of Mazandaran University of Medical Sciences (JMUMS)*. 2014;24(119):213-22.
- [16] Mahmoudi R, Asadpour R, Pajouhi Alamoti MR, Golchin A, Kiyani R, Mohammadpour R. Raw cow milk quality: Relationship between antibiotic residue and Somatic cell count. *Int F Res J*. 2013;20(6):3347-50.
- [17] Mahmoudi R, Amini K, Vagef R, Vahab M, Mir H, Vagef R. Antibiotics residues in raw and pasteurized milk, Iran. *J of Res & Heal*. 2014;4(4):884-89.
- [18] Manafi M, Hesaraki J, Rafat SA. Monitoring of antibiotic residue in raw and pasteurised milk in East Azerbaijan of Iran by delvotest method. *J Food Research Agricultural Science*. 2011;20(3):125-31.
- [19] Bahreinipour A, Mohsenzadeh M. Identification of antimicrobial residual compounds in raw and pasteurized milk by Yoghurt Culture Test. *Sci-Res Iran Vat J*. 2009;3(24):5-11.
- [20] Ghanavi Z, Mollayi S, Eslami Z. Determination between the amount of Penicillin G residue in raw and pasteurized milk in Iran. *Jin J Mic*. 2013;6(7):e12724.
- [21] Abedi Shirazi Kh. Survey on the contamination of milk with antibiotics in Shiraz. (Thesis). *Fac of Vet Med Uni of Shir*. 1983.
- [22] Lebreton MH, Savoy-Perroud MC, Diserens JM. Validation and comparison of the Copan Milk Test and delvotest SP-NT for the detection of antimicrobials in milk. *Ana Chim Acta*. 2006;586(1):280-83.
- [23] Bilandzic N, Solomun Kolanovic B, Varenina I, Scortichini G, Annunziata L, Brstilo M, et al. Veterinary drug residues determination in raw milk in Croatia. *F Con*. 2011;22:1941-48.
- [24] Rassouli A, Faghghi SM. A survey on antibacterial drug usage in dairy farms in Tehran province during 2004-2005. *J of Vet Res*. 2014;69(1):41-48.
- [25] Rassouli A, Abdolmaleki Z, Bokaei S, Kamkar A, Shams G. A cross sectional study on oxytetracycline and tetracycline residues in pasteurized milk supplied in Tehran by an HPLC method. *Int J Vet Res*. 2010;4(1):1-3.
- [26] Fallahrad AH, Mohsenzadeh M, Asadpour HR. Determination of gentamycin residue in raw milk delivered to the milk factory of Mashad. *Sci and Agric* 2005;20(7):183-89. [In Persian].
- [27] Dabbagh Moghaddam A, Tayebi L, Falahatpisheh H, Mahmoudian M, Kowsari N, Akbarein H, et al. Evaluation of the tetracycline residues in pasteurized milks distributed in Tehran by HPLC method. *J Army Univ Med Sci*. 2014;11(4):318-23. [In Persian].
- [28] Mohammad Sadegh M, Bokaei S, Moosavi T, Abbaszade E. Comparing the efficacy of Copan and Delvo test to detect of antibiotic residues in bulk tank milk samples of Garmsar dairy farms. *J of Vet Micr*. 2007;3(1):23-28.
- [29] Mohamadi Sani A, Nikpooyan H, Moshiri R. Aflatoxin M1 contamination and antibiotic residue in milk in Khorasan province. Iran. *F and Chem Tox*. 2010;48:2130-32.
- [30] Yamaki M, Berruga MI, Althaus RL, Molina MP, Molina A. Occurrence of antibiotic residues in milk from Manchega ewe dairy farms. *J of Dairy Sci*. 2004;87: 3132-37.
- [31] Ghidini S, Zanardi E, Varisco G, Chizzolini R. Prevalence of molecules of β -lactam antibiotics in bovine milk in Lombardia and Emilia Romagna (Italy). *Ann Fac MedicVet Di Parma*. 2002;22:245-52.

PARTICULARS OF CONTRIBUTORS:

1. PhD. Student, Deputy of Food and Drugs, Gonabad University of Medical Sciences, Gonabad, I.R, Iran.
2. Assistant Professor in TEFL, Department of Basic Sciences, Faculty of Medicine, Gonabad University of Medical Sciences, Gonabad, Iran.
3. Faculty of Health, Department of Public Health, Gonabad University of Medical Sciences, Gonabad, I.R, Iran.
4. Student, Department of Basic Sciences, Gonabad University of Medical Sciences, Gonabad, I.R, Iran.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Hamed Ramezani Awal Riabi,
Faculty of Health, Department of Public Health, Gonabad University of Medical Sciences, Gonabad, I.R, Iran.
E-mail: hamedramazany@yahoo.com

Date of Submission: **May 01, 2016**
Date of Peer Review: **Jun 08, 2016**
Date of Acceptance: **Aug 29, 2016**
Date of Publishing: **Dec 01, 2016**

FINANCIAL OR OTHER COMPETING INTERESTS: None.